

TOPOLOGY OF WORDS

VLADIMIR TURAEV

ABSTRACT. We introduce a topological approach to words. Words are approximated by Gauss words and then studied up to natural modifications inspired by homotopy transformations of curves on the plane.

1. INTRODUCTION

Words are finite sequences of letters in a given alphabet. Every word has its own personality and should be treated with the same respect and attention as say, a polyhedron or a manifold. In this paper we attempt to study words as topological objects. A word in an alphabet α can be viewed as a way of interaction or interlacement of the letters of α . For example, two letters $a, b \in \alpha$ are interlaced in the word $abab$ and are not interlaced in the word $aabb$. From this perspective, a word can be compared with a link of circles in Euclidean 3-space, the letters being the counterparts of the circles.

Another geometric viewpoint is suggested by the Gauss-Rosenstiehl [Ro] correspondence between certain words and plane curves. This viewpoint consists in treating an arbitrary word in the alphabet α as a “curve” passing through the points of α in the prescribed order. To make this work, one needs to embed α in a bigger space, say a surface, or better to label certain points of a surface with letters of α . These ideas, albeit imprecise at this stage, suggest further directions of thought. First of all, the letters appearing in the word with multiplicity ≥ 3 are “singular” self-crossings of the curve. They may be “desingularized” to obtain only double self-crossings, see Figure 1. Secondly, one may allow different points of the ambient surface to be labeled with the same letter which leads us to so-called étale words generalizing the ordinary words. Thirdly, the notion of homotopy for curves leads us to a notion of homotopy for words.

The key new concepts introduced in this paper are those of étale words and nanowords. An étale word over an alphabet α is a word in an alphabet $\mathcal{A}$ endowed




FIGURE 1. Desingularization




FIGURE 2. Local deformations of plane curves

with a projection $\mathcal{A} \rightarrow \alpha$. The image of a letter $A \in \mathcal{A}$ under this projection is denoted $|A|$. For instance, $ABCABC$ is an étale word over α provided we specify $|A|, |B|, |C| \in \alpha$. Every word in the alphabet α becomes an étale word over α by using the identity $\text{id} : \alpha \rightarrow \alpha$ as the projection. An étale word in which every letter appears twice or not at all is called a nanoword. For instance, $ABCABC$ is a nanoword. Every étale word can be approximated by a nanoword via a desingularization which replaces each letter of multiplicity $m \geq 3$ by $m(m-1)/2$ letters of multiplicity 2. This allows us to focus on nanowords; all definitions and results concerning them extend to arbitrary étale words (and, in particular, to arbitrary words) via desingularization.

We now fix an additional piece of data: an involution $\tau : \alpha \rightarrow \alpha$ (it may be the identity). Given τ , we introduce an equivalence relation of homotopy on the set of étale words over α . It is implied that homotopic étale words give rise to the same interlacement of letters of α with respect to τ . The relation of homotopy is generated by three transformations or moves on nanowords. The first move consists in deleting two consecutive entries of the same letter. The second move has the form $xAB_yBAz \mapsto xyz$ where x, y, z are words and A, B are letters such that $\tau(|A|) = |B|$. The third move has the form $xAB_yAC_zBCt \mapsto xBA_yCA_zCBt$ where x, y, z, t are words and A, B, C are letters such that $|A| = |B| = |C|$. These moves are suggested by the standard local deformations of plane curves, see Figure 2.

We shall study properties and characteristics of words and étale words preserved under homotopy. A number of methods developed in topology find an echo in this setting. The homological intersection theory of curves on surfaces suggests a family of homotopy invariants of nanowords (the self-linking function, the linking form, the linking pairing). The technique of Alexander matrices in knot theory leads us to modules and polynomial invariants of nanowords including a rather powerful invariant λ . The technique of colorings of knot diagrams suggests a notion of a coloring for nanowords. The theory of knot quandles finds its analogue in the form of α -keis. The theory of virtual strings leads to so-called characteristic sequences of words. Other methods of low-dimensional topology may possibly apply in this setting.

As an application of our algebraic methods, we give a homotopy classification of nanowords of length ≤ 6 . All nanowords of length 2 are contractible, i.e., homotopic to an empty nanoword. The classification of nanowords of length 4 is as follows: the nanowords of the form $AABB$ or $ABBA$ are contractible (by the first move); a nanoword of the form $ABAB$ is contractible if and only if $|A| = \tau(|B|)$;

two non-contractible nanowords $ABAB$ and $A'B'A'B'$ are homotopic if and only if $|A| = |A'|$ and $|B| = |B'|$ (Theorem 8.4.1). The homotopy classification of nanowords of length 6 is more involved. Excluding those homotopic to nanowords of length ≤ 4 by the first move, we obtain 5 families of nanowords of the form

$$ABCABC, ABCACB, ABCBAC, ABCBCA, ABACBC.$$

We fully describe when such nanowords are homotopic to nanowords of length ≤ 4 and when they are homotopic to each other (Theorem 13.2.1).

We also give a homotopy classification of words of length ≤ 5 in the alphabet α (Theorems 9.0.1 and 9.0.2). For $\tau = \text{id} : \alpha \rightarrow \alpha$ this classification is especially simple: a word of length ≤ 5 , in which every letter appears at least twice or not at all, is non-contractible if and only if it has one of the following six forms: $abab, abaab, baaba, aabab, babaa, ababa$ with distinct $a, b \in \alpha$. Two words from this list are homotopic if and only if they coincide letter-wise.

In the case where the alphabet α consists of two elements permuted by τ , the notion of a nanoword over α is equivalent to the notion of an open virtual string introduced in [Tu1]. The results of the present paper generalize a number of results of [Tu1] and of the subsequent paper by D. Silver and S. Williams [SW] on open virtual strings. In particular, the invariant λ and the characteristic sequences generalize the invariants of open strings introduced by Silver and Williams.

To make this paper accessible to readers not interested in topology, it is written in purely algebraic terms. Relations with topology will be discussed elsewhere.

The paper consists of 4 parts. The first part (Sections 2 – 5) is devoted to the basics. We define étale words, nanowords, desingularization of étale words, homotopy and coverings of nanowords, and discuss a group-theoretic approach to nanowords. In the second part (Sections 6 – 9) we construct linking invariants of nanowords and give a homotopy classification of nanowords of length 4 and of words of length ≤ 5 . The third part (Sections 10 – 13) begins with a discussion of colorings of nanowords and proceeds to modules and polynomials of nanowords. Then we give a homotopy classification of nanowords of length 6. In the fourth part (Sections 14 – 16) we introduce α -keis and characteristic sequences of nanowords. They are used to distinguish certain nanowords of length 6 from nanowords of length 4 and to accomplish thus the homotopy classification of nanowords of length ≤ 6 .

Throughout the paper the symbol α denotes a set endowed with an involution $\tau : \alpha \rightarrow \alpha$.

2. ÉTALE WORDS AND NANOWORDS

2.1. Words. An *alphabet* is a set and *letters* are its elements. A *word of length* $n \geq 1$ in an alphabet α is a mapping $w : \hat{n} \rightarrow \alpha$ where $\hat{n} = \{1, 2, \dots, n\}$. Such a word is encoded by the sequence $w(1)w(2)\cdots w(n)$. For example, the sequence aba in the alphabet $\alpha = \{a, b\}$ encodes the word $\hat{3} \rightarrow \alpha$ sending 1, 2, 3 to a, b, a respectively. By definition, there is a unique *empty word* $\emptyset$ of length 0.

Writing down consecutively the letters of two words w and v we obtain their concatenation wv . For instance, the concatenation of $w = abb$ and $v = aa$ is the

word $wv = abbaa$. Writing the letters of a word w in the opposite order we obtain the *opposite word* w^- . For instance, if $w = abb$, then $w^- = bba$.

The *multiplicity* of a letter $a \in \alpha$ in a word $w : \hat{n} \rightarrow \alpha$ is the number $m = m_w(a) = \text{card}\{i \in \hat{n} \mid w(i) = a\}$. We say that a appears m times in w . For example, $m_{aab}(a) = 2$ and $m_{aab}(b) = 1$.

A *monoliteral* word of length $m \geq 2$ is the word $aa \cdots a$ formed by m copies of the same letter $a \in \alpha$. This word is denoted a^m .

A mapping f from a set α_1 to a set α_2 induces a mapping $f_\#$ from the set of words in the alphabet α_1 to the set of words in the alphabet α_2 . It is obtained by applying f letterwise.

2.2. Étale words. The class of words in the alphabet α is too narrow for our purposes. We introduce here a wider class of étale words over α .

An α -*alphabet* is a set $\mathcal{A}$ endowed with a mapping $\mathcal{A} \rightarrow \alpha$ called *projection*. The image of a letter $A \in \mathcal{A}$ under the projection is denoted $|A|$. Any subset $\mathcal{A}'$ of an α -alphabet $\mathcal{A}$ becomes an α -alphabet by restricting the projection $\mathcal{A} \rightarrow \alpha$ to $\mathcal{A}'$.

A *morphism* of α -alphabets $\mathcal{A}_1, \mathcal{A}_2$ is a set-theoretic mapping $f : \mathcal{A}_1 \rightarrow \mathcal{A}_2$ such that $|A| = |f(A)|$ for all $A \in \mathcal{A}_1$. If f is bijective, then this morphism is an *isomorphism*.

An *étale word* over α is a pair (an α -alphabet $\mathcal{A}$, a word in the alphabet $\mathcal{A}$). Two étale words $(\mathcal{A}_1, w_1)$ and $(\mathcal{A}_2, w_2)$ over α are *isomorphic* if there is an isomorphism $f : \mathcal{A}_1 \rightarrow \mathcal{A}_2$ such that $w_2 = f_\#(w_1)$. The relation of isomorphism for étale words is denoted $\approx$.

We define a product of étale word $(\mathcal{A}_1, w_1)$ and $(\mathcal{A}_2, w_2)$ over α as follows. Replacing if necessary $(\mathcal{A}_1, w_1)$ with an isomorphic étale word we can assume that $\mathcal{A}_1 \cap \mathcal{A}_2 = \emptyset$. Then the product in question is the étale word $(\mathcal{A}_1 \cup \mathcal{A}_2, w_1 w_2)$ over α . It is well defined up to isomorphism. Multiplication of étale words is associative and has a unit represented by an empty étale word in an empty α -alphabet.

For each étale word $(\mathcal{A}, w)$, we have the *opposite* étale word $(\mathcal{A}, w^-)$ where w^- is the word opposite to w . For each α -alphabet $(\mathcal{A}, p : \mathcal{A} \rightarrow \alpha)$, the *inverse* α -alphabet $\overline{\mathcal{A}}$ is the same set $\mathcal{A}$ with projection $\tau p : \mathcal{A} \rightarrow \alpha$ where $\tau : \alpha \rightarrow \alpha$ is the fixed involution. The *inverse* of an étale word $(\mathcal{A}, w)$ over α is the étale word $\overline{w} = (\overline{\mathcal{A}}, w)$. An étale word w is *symmetric* (resp. *skew-symmetric*) if it is isomorphic to w^- (resp. to $\overline{w}^-$). For instance, vv^- is symmetric and $v\overline{v}^-$ is skew-symmetric for any étale word v .

Each word w in the alphabet α gives rise to an étale word $(\mathcal{A} = \alpha, w)$ over α where the projection $\mathcal{A} \rightarrow \alpha$ is the identity. In this way, étale words over α generalize words in the alphabet α .

Warning: concatenation of words in the alphabet α differs from multiplication of the corresponding étale words. For instance, for the words abb and aa in the alphabet $\alpha = \{a, b\}$, the corresponding étale words are $(\mathcal{A}_1 = \{A, B\}, ABB)$ with $|A| = a, |B| = b$ and $(\mathcal{A}_2 = \{C\}, CC)$ with $|C| = a$. Their product is the étale word $(\{A, B, C\}, ABBC C)$ with $|A| = |C| = a, |B| = b$. On the other hand, the étale word corresponding to $abbaa$ is $(\{A, B\}, ABBA A)$ with $|A| = a, |B| = b$.

2.3. Gauss words and nanowords. A word w in a finite alphabet $\mathcal{A}$ is a *Gauss word* if every letter of $\mathcal{A}$ appears in w exactly twice. For instance, $ABAB$ is a Gauss word in the alphabet $\{A, B\}$ while ABA and AA are not Gauss words in this alphabet. If $\mathcal{A}$ consists of N letters, then there are $(2N)!$ Gauss words in the alphabet $\mathcal{A}$. Note that concatenation of two or more Gauss words in a non-empty alphabet is never a Gauss word.

A *nanoword* is an étale Gauss word. More precisely, a *nanoword over α* is a pair (a finite α -alphabet $\mathcal{A}$, a Gauss word in the alphabet $\mathcal{A}$). The *length* of this nanoword is the length of the Gauss word in question, i.e., $2 \operatorname{card}(\mathcal{A})$.

The set of nanowords over α is denoted $\mathcal{N}(\alpha)$. Two nanowords are *isomorphic* if they are isomorphic as étale words. Observe that the étale word opposite or inverse to a nanoword is a nanoword. The product of two nanowords is a nanoword (defined up to isomorphism). An empty étale word in an empty α -alphabet is a nanoword called the *empty nanoword*.

Instead of writing $(\mathcal{A}, w)$ for a nanoword over α , we shall often write simply w . The alphabet $\mathcal{A}$ can be uniquely recovered from w as the set of all letters appearing in w . However, the projection $\mathcal{A} \rightarrow \alpha$ should be always specified.

Each Gauss word w in a finite alphabet α gives rise to a nanoword $(\mathcal{A} = \alpha, w)$ over α where the projection $\mathcal{A} \rightarrow \alpha$ is the identity. In this way, nanowords over α generalize Gauss words in the alphabet α .

2.4. Example. Let $\alpha = \{a, b\}$ with involution permuting a and b . Let $\mathcal{A} = \{A_1, A_2, B_1\}$ with $|A_1| = |A_2| = a, |B_1| = b$. Then $(\mathcal{A}, A_1 B_1 A_1 A_2 A_2 B_1)$ is a nanoword over α . The opposite nanoword is $(\mathcal{A}, B_1 A_2 A_2 A_1 B_1 A_1)$ and the inverse is $(\overline{\mathcal{A}}, A_1 B_1 A_1 A_2 A_2 B_1)$ where $\overline{\mathcal{A}} = \{A_1, A_2, B_1\}$ with $|A_1| = |A_2| = b, |B_1| = a$.

2.5. Desingularization. For an étale word $(\mathcal{A}, w)$ over α , we define a nanoword $(\mathcal{A}^d, w^d)$ over α called the *desingularization* of w . The alphabet $\mathcal{A}^d$ consists of the triples (A, i, j) where $A \in \mathcal{A}$ and $1 \leq i < j \leq m_w(A)$ where $m_w(A)$ is the multiplicity of A in w . For brevity, we write $A_{i,j}$ for (A, i, j) . We make $\mathcal{A}^d$ into an α -alphabet by $|A_{i,j}| = |A| \in \alpha$ for all A, i, j . The word w^d is obtained from w by first deleting all $A \in \mathcal{A}$ with $m_w(A) = 1$. Then for each $A \in \mathcal{A}$ with $m_w(A) \geq 2$ and each $i = 1, 2, \dots, m_w(A)$, we replace the i -th entry of A in w by

$$A_{1,i} A_{2,i} \cdots A_{i-1,i} A_{i,i+1} A_{i,i+2} \cdots A_{i,m_w(A)}.$$

The resulting word w^d in the alphabet $\mathcal{A}^d$ is a Gauss word so that $(\mathcal{A}^d, w^d)$ is a nanoword of length $\sum_{A \in \mathcal{A}} m_w(A)(m_w(A) - 1)$. For example, if $w = AABABC$, then $\mathcal{A}^d = \{A_{1,2}, A_{1,3}, A_{2,3}, B_{1,2}\}$ and $w^d = A_{1,2} A_{1,3} A_{1,2} A_{2,3} B_{1,2} A_{1,3} A_{2,3} B_{1,2}$.

The desingularization may considerably increase the length of a word but this is the price to pay for obtaining a Gauss word. If w is a nanoword, then $w^d \approx w$. If $w = \emptyset$, then $w^d = \emptyset$. It follows from the definitions that $(w_1 w_2)^d = w_1^d w_2^d$ for any étale words w_1, w_2 , $\overline{w}^d = \overline{w^d}$, and $(w^-)^d \approx (w^d)^-$ for any w . (The latter isomorphism sends $A_{i,j}$ to $A_{m_w(A)+1-j, m_w(A)+1-i}$ for all A, i, j).

3. HOMOTOPY

3.1. Homotopy moves. To define homotopy of nanowords we fix a *homotopy data* consisting of a set α with involution $\tau : \alpha \rightarrow \alpha$ and a set $S \subset \alpha^3 = \alpha \times \alpha \times \alpha$. Given this data, we define three transformations of nanowords over α called *S-homotopy moves* or, when S is fixed, *homotopy moves*.

(1). The first move applies to any nanoword of the form $(\mathcal{A}, xAAy)$ where $A \in \mathcal{A}$ and x, y are words in the alphabet $\mathcal{A}' = \mathcal{A} - \{A\}$. It transforms $(\mathcal{A}, xAAy)$ into the nanoword $(\mathcal{A}', xy)$ where the structure of an α -alphabet in $\mathcal{A}'$ is obtained by restricting the one in $\mathcal{A}$. Note that xy is a Gauss word in the alphabet $\mathcal{A}'$.

The inverse move $(\mathcal{A}', xy) \mapsto (\mathcal{A}' \cup \{A\}, xAAy)$ adds a new letter A to $\mathcal{A}'$ with arbitrary $|A| \in \alpha$ and replaces the Gauss word xy in the α -alphabet $\mathcal{A}'$ with $xAAy$.

(2). The second move applies to a nanoword of the form $(\mathcal{A}, xAByBAz)$ where $A, B \in \mathcal{A}$ with $|B| = \tau(|A|)$ and x, y, z are words in the alphabet $\mathcal{A}' = \mathcal{A} - \{A, B\}$. This nanoword is transformed into $(\mathcal{A}', xyz)$ where the structure of an α -alphabet in $\mathcal{A}'$ is obtained by restricting the one in $\mathcal{A}$.

The inverse move $(\mathcal{A}', xyz) \mapsto (\mathcal{A}' \cup \{A, B\}, xAByBAz)$ adds two new letters A, B to $\mathcal{A}'$ with arbitrary $|A| \in \alpha$ and $|B| = \tau(|A|)$ and replaces the Gauss word xyz in the alphabet $\mathcal{A}'$ with $xAByBAz$.

(3) The third move applies to a nanoword of the form $(\mathcal{A}, xAByACzBCt)$ where $A, B, C \in \mathcal{A}$ are distinct letters with $(|A|, |B|, |C|) \in S$ and x, y, z, t are words in the alphabet $\mathcal{A} - \{A, B, C\}$ such that $xyzt$ is a Gauss word in this alphabet. The move transforms $(\mathcal{A}, xAByACzBCt)$ into $(\mathcal{A}, xBAyCAzCBt)$. The inverse move transforms $(\mathcal{A}, xBAyCAzCBt)$ into $(\mathcal{A}, xAByACzBCt)$. It applies if $(|A|, |B|, |C|) \in S$.

3.2. Homotopy of nanowords. *Homotopy* is the equivalence relation in the class of nanowords generated by isomorphism and the homotopy moves. For a homotopy data (α, S) , two nanowords over α are *S-homotopic* if they can be obtained from each other by a finite sequence of *S-homotopy moves* (1) – (3), the inverse moves, and isomorphisms. The relation of *S-homotopy* is denoted $\simeq_S$. A nanoword *S-homotopic* to an empty nanoword $\emptyset$ is said to be *S-contractible*. All these notions crucially depend on the choice of $\tau : \alpha \rightarrow \alpha$ and S .

The set of *S-homotopy* classes of nanowords over α is denoted $\mathcal{N}_\bullet^S(\alpha) = \mathcal{N}(\alpha) / \simeq_S$. Multiplication of nanowords is compatible with *S-homotopy*: if nanowords w_1, w_2 are *S-homotopic* to nanowords w'_1, w'_2 , respectively, then $w_1 w_2$ is *S-homotopic* to $w'_1 w'_2$. Multiplication of nanowords makes $\mathcal{N}_\bullet^S(\alpha)$ into a monoid with unit represented by $\emptyset$. If S is invariant under the involution $\tau \times \tau \times \tau$ on α^3 , then nanowords inverse to *S-homotopic* nanowords are themselves *S-homotopic*. If S is invariant under the involution $(a, b, c) \mapsto (c, b, a)$ on α^3 , then nanowords opposite to *S-homotopic* nanowords are *S-homotopic*.

A nanoword w is *S-homotopically symmetric* (resp. *S-homotopically skew-symmetric*) if $w \simeq_S w^-$ (resp. if $w \simeq_S \overline{w}^-$). Clearly, (skew-) symmetric nanowords are *S-homotopically* (skew-) symmetric. *S-contractible* nanowords are *S-homotopically* symmetric and skew-symmetric. If nanowords w, w' are *S-homotopic*,

then $w'w^-$ is S -homotopically symmetric and $w'\overline{w}^-$ is S -homotopically skew-symmetric.

For a nanoword w , let $\|w\|_S$ be half of the minimal length of a nanoword homotopic to w . This defines a $\mathbb{Z}$ -valued “norm” $\|\cdot\|_S$ on $\mathcal{N}_\bullet^S(\alpha)$ whose value on an S -homotopy class of nanowords is half of the minimal length of a nanoword in this class. Clearly, $\|w\|_S = 0$ if and only if w is S -contractible. Since all nanowords of length 2 are S -contractible, the function $\|\cdot\|_S$ does not take the value 1. It is obvious that $\|w_1w_2\|_S \leq \|w_1\|_S + \|w_2\|_S$ for any w_1, w_2 .

Lemma 3.2.1. *Let A, B, C be distinct letters in an α -alphabet $\mathcal{A}$ and let x, y, z, t be words in the alphabet $\mathcal{A} - \{A, B, C\}$ such that $xyzt$ is a Gauss word in this alphabet. Then*

- (i) $(\mathcal{A}, xAB yCAzBCt) \simeq_S (\mathcal{A}, xBA yACzCBt)$ if $(|A|, \tau(|B|), |C|) \in S$,
- (ii) $(\mathcal{A}, xAB yCAzCBt) \simeq_S (\mathcal{A}, xBA yACzBCt)$ if $(\tau(|A|), \tau(|B|), |C|) \in S$,
- (iii) $(\mathcal{A}, xAB yACzCBt) \simeq_S (\mathcal{A}, xBA yCAzBCt)$ if $(|A|, \tau(|B|), \tau(|C|)) \in S$.

Proof. Pick two symbols D, E not belonging to $\mathcal{A}$.

We verify (i). Set $|D| = |B| \in \alpha$ and $|E| = \tau(|B|) \in \alpha$. Applying an inverse (2)-move inserting $\cdots DE \cdots ED \cdots$ and an inverse (3)-move we obtain

$$\begin{aligned} (\mathcal{A}, xAB yCAzBCt) &\simeq_S (\mathcal{A} \cup \{D, E\}, xD \underline{E} A B y \underline{C} A z B \underline{C} E D t) \\ &\simeq_S (\mathcal{A} \cup \{D, E\}, xDA E B y A C z B E C D t) \end{aligned}$$

where for visual convenience the letters modified by the inverse (3)-move are underlined. Here we use that $(|A|, |E|, |C|) \in S$. Applying to the resulting nanoword a (2)-move removing B, E we obtain the nanoword $(\mathcal{A} \cup \{D\} - \{B\}, xDA y A C z C D t)$. It is isomorphic to $(\mathcal{A}, xBA y A C z C B t)$.

We verify (ii). Set $|D| = \tau(|A|)$ and $|E| = |A|$. Applying an inverse (2)-move inserting $\cdots DE \cdots ED \cdots$ and an inverse to the homotopy (i) we obtain

$$\begin{aligned} (\mathcal{A}, xAB yCAzCBt) &\simeq_S (\mathcal{A} \cup \{D, E\}, x \underline{A} B \underline{D} E y E \underline{D} C A z \underline{C} B t) \\ &\simeq_S (\mathcal{A} \cup \{D, E\}, x \underline{A} D B E y E C \underline{D} A z B C t) \simeq_S (\mathcal{A} \cup \{E\} - \{A\}, xB E y E C z B C t). \end{aligned}$$

The latter nanoword is isomorphic to $(\mathcal{A}, xBA y A C z B C t)$.

To prove (iii), set $|D| = |C|$ and $|E| = \tau(|C|)$. Then

$$\begin{aligned} xAB yACzCBt &\simeq_S x \underline{A} B y D E A C z C B E D t \simeq_S xBA yDA E C z C E B D t \\ &\simeq_S xBA yDA zBDt \approx xBA yCAzBCt. \end{aligned}$$

□

Lemma 3.2.2. *Suppose that $S \cap (\alpha \times b \times b) \neq \emptyset$ for all $b \in \alpha$. Let $(\mathcal{A}, xAB yABz)$ be a nanoword over α where $A, B \in \mathcal{A}$ with $|B| = \tau(|A|)$ and x, y, z are words in $\mathcal{A} - \{A, B\}$. Then $(\mathcal{A}, xAB yABz) \simeq_S (\mathcal{A} - \{A, B\}, xyz)$.*

Proof. Set $b = |B| \in \alpha$. By assumption, there is $e \in \alpha$ such that $(e, b, b) \in S$. Pick a symbol E not belonging to $\mathcal{A}$ and set $|E| = \tau(e)$. Then

$$\begin{aligned} (\mathcal{A}, xAB yABz) &\simeq_S (\mathcal{A} \cup \{E\}, x\underline{AE} \underline{EB} y \underline{AB} z) \simeq_S (\mathcal{A} \cup \{E\}, xE \underline{ABE} y \underline{BA} z) \\ &\simeq_S (\mathcal{A} \cup \{E\} - \{A, B\}, xEEyz) \simeq_S (\mathcal{A} - \{A, B\}, xyz). \end{aligned}$$

Here the second homotopy is provided by item (ii) of the previous lemma where A, B, C are replaced with E, A, B , respectively. We use that $(\tau(|E|), \tau(|A|), |B|) = (e, b, b) \in S$. $\square$

3.3. Homotopy of étale words. We say that étale words w_1, w_2 over α are S -homotopic and write $w_1 \simeq_S w_2$ if $w_1^d \simeq_S w_2^d$. Clearly, isomorphic étale words are S -homotopic. S -homotopy is an equivalence relation in the class of étale words. For nanowords, this relation of S -homotopy coincides with the one defined above. Every S -homotopy invariant I of nanowords extends to an S -homotopy invariant I^d of étale words by $I^d(w) = I(w^d)$. In particular, the S -norm of an étale word w is defined by $\|w\|_S = \|w^d\|_S$.

An étale word w is S -contractible (resp. S -homotopically (skew-) symmetric) if the nanoword w^d is S -contractible (resp. S -homotopically (skew-) symmetric). Since words in the alphabet α may be treated as étale words, we can apply to them all these definitions.

Although there are various interesting choices for S , we focus here on one simplest S . In the remaining part of the paper we assume that S is the diagonal of α^3 that is $S = \{(a, a, a)\}_{a \in \alpha}$. In other words we allow the third homotopy move $(\mathcal{A}, xAB yAC zBCt) \mapsto (\mathcal{A}, xBA yCA zCBt)$ and the inverse move if and only if $|A| = |B| = |C|$. Under this convention, we shall omit the prefix S - and speak simply of homotopy rather than S -homotopy. We shall also omit the index S and write $\simeq, \|\cdot\|, \mathcal{N}_\bullet(\alpha)$ for $\simeq_S, \|\cdot\|_S, \mathcal{N}_\bullet^S(\alpha)$. Note that $S \cap (\alpha \times b \times b) = (b, b, b)$ for $b \in \alpha$ so that we can apply Lemma 3.2.2.

The principal aim of the homotopy theory of nanowords is a computation of $\mathcal{N}_\bullet(\alpha)$. As a step in this direction, we shall construct several homotopy invariants of nanowords. They will often allow us to distinguish nanowords up to homotopy and to decide whether a given nanoword is homotopically (skew-) symmetric or homotopic to its inverse. A related problem is a computation or at least an estimate for $\|\cdot\|$.

The following lemma shows that for words in a given alphabet, the relation of homotopy does not change under a passage to a bigger alphabet.

Lemma 3.3.1. *Let β be a τ -invariant subset of α . If two étale words over β are homotopic in the class of étale words over α , then they are homotopic in the class of étale words over β .*

Proof. Given an étale word $(\mathcal{A}, w)$ over α we define its *pull-back* to β to be the étale word over β obtained by deleting from both $\mathcal{A}$ and w all letters $A \in \mathcal{A}$ with $|A| \in \alpha - \beta$. The pull-back commutes with desingularization and transforms isomorphic (resp. homotopic) étale words over α into isomorphic (resp. homotopic) étale words over β .

Observe that any equivariant mapping of sets with involution $f : \alpha' \rightarrow \alpha$ induces a mapping $f_{\#} : \mathcal{N}(\alpha') \rightarrow \mathcal{N}(\alpha)$ by $f_{\#}(\mathcal{A}, p : \mathcal{A} \rightarrow \alpha', w) = (\mathcal{A}, fp : \mathcal{A} \rightarrow \alpha, w)$. Clearly, $f_{\#}$ transforms isomorphic (resp. homotopic) nanowords into isomorphic (resp. homotopic) nanowords. Denote the induced mapping $\mathcal{N}_{\bullet}(\alpha') \rightarrow \mathcal{N}_{\bullet}(\alpha)$ by $f_{\bullet}$. The composition of $f_{\bullet}$ with the pull-back to $f(\alpha') \subset \alpha$ is the mapping $\mathcal{N}_{\bullet}(\alpha') \rightarrow \mathcal{N}_{\bullet}(f(\alpha))$ induced by f . If f is injective, then the latter mapping is an isomorphism and therefore $f_{\bullet}$ is injective. Applying this to the inclusion $\beta \hookrightarrow \alpha$, we obtain the claim of the lemma. $\square$

3.4. Examples. 1. The nanoword $(\mathcal{A} = \{A\}, AA)$ is contractible for any choice of $|A| \in \alpha$. The nanowords $ABBA$ and $AABB$ (in the alphabet $\{A, B\}$) are contractible for any $|A|, |B| \in \alpha$. The nanoword $ABAB$ is contractible provided $|A| = \tau(|B|)$. The nanoword $ABCDABCD$ (in the alphabet $\{A, B, C, D\}$) is contractible provided either $|A| = \tau(|B|)$ and $|C| = \tau(|D|)$ or $|B| = \tau(|C|)$ and $|A| = \tau(|D|)$.

2. A more sophisticated example: the nanoword $ABCDEFBGDHFAGCHE$ is contractible provided $|A| = |B| = |F| = \tau(|C|) = \tau(|D|) = \tau(|G|)$ (the elements $|E|, |H| \in \alpha$ may be arbitrary). Indeed, applying Lemma 3.2.1(iii), Lemma 3.2.2, and the first homotopy move we obtain

$$\begin{aligned} \underline{ABCDEFBGDHFAGCHE} &\simeq \underline{ACBDEFGBDHFACGHE} \\ &\simeq \underline{BDEFGBDHFEGHE} \simeq \underline{BDEBDHHE} \simeq \underline{EHHE} \simeq \underline{EE} \simeq \emptyset. \end{aligned}$$

Note that there is no way to apply to the nanoword $ABCDEFBGDHFAGCHE$ the first or second homotopy moves; to contract this nanoword one needs the third homotopy move.

3. Let $a \in \alpha$ be a letter such that $\tau(a) = a$. We show that the monoliteral words a^3, a^4, a^5 are contractible. The étale word corresponding to $a^3 = aaa$ is $(\mathcal{A} = \alpha, aaa)$ and the desingularization yields the nanoword $-12-13-12-23-13-23$ where we write $-ij$ for $a_{i,j}$. We have

$$\underline{-12-13-12-23-13-23} \simeq \underline{-13-12-23-12-23-13} \simeq \underline{-13-13} \simeq \emptyset.$$

Here is a contracting homotopy for the desingularization of $a^4 = aaaa$:

$$\begin{aligned} &\underline{-12-13-14-12-23-24-13-23-34-14-24-34} \\ &\simeq \underline{-13-12-14-23-12-24-23-13-34-14-24-34} \\ &\simeq \underline{-13-14-12-23-24-12-23-13-34-24-14-34} \\ &\simeq \underline{-13-14-24-13-34-24-14-34} \simeq \underline{-13-13-34-34} \simeq \emptyset. \end{aligned}$$

Here is a contracting homotopy for the desingularization of $a^5 = aaaaa$:

$$\begin{aligned} &\underline{-12-13-14-15-12-23-24-25-13-23-34-35-14-24-34-45-15-25-35-45} \\ &\simeq \underline{-13-12-14-15-23-12-24-25-23-13-34-35-14-24-34-45-15-25-35-45} \\ &\simeq \underline{-13-14-12-15-23-24-12-25-23-13-34-35-24-14-34-45-15-25-35-45} \\ &\simeq \underline{-13-14-15-12-23-24-25-12-23-13-34-35-24-14-34-45-25-15-35-45} \\ &\simeq \underline{-13-14-15-24-25-13-34-35-24-14-34-45-25-15-35-45} \\ &\simeq \underline{-14-13-15-24-25-34-13-35-24-34-14-45-25-15-35-45} \end{aligned}$$

$$\begin{aligned}
& \simeq -14-15-13-24-25-34-35-13-24-34-14-45-25-35-15-45 \\
& \simeq -14-15-25-34-35-34-14-45-25-35-15-45 \\
& \simeq -15-14-25-34-35-34-45-14-25-35-45-15 \\
& \simeq -15-34-35-34-45-35-45-15 \\
& \simeq -15-35-34-45-34-45-35-15 \simeq -34-45-34-45 \simeq \emptyset.
\end{aligned}$$

Using a topological argument one can show that a^m is contractible for all $m \geq 2$. (One can realize the desingularization of a^m as the Gauss word of a generic closed curve in $\mathbf{R}^2$ and use that all curves in $\mathbf{R}^2$ are contractible).

4. Let $a \in \alpha$ be a letter such that $\tau(a) \neq a$. Set $b = \tau(a)$. Then the word $w = abab$ is contractible. Indeed, the desingularization yields the nanoword

$$(\mathcal{A}^d = \{a_{1,2}, a_{1,3}, a_{2,3}, b_{1,2}\}, w^d = a_{1,2}a_{1,3}b_{1,2}a_{1,2}a_{2,3}b_{1,2}a_{1,3}a_{2,3})$$

where $|a_{i,j}| = a$ for all i, j and $|b_{1,2}| = b$. We have

$$\begin{aligned}
w^d &= \underline{a_{1,2}a_{1,3}} \underline{b_{1,2}a_{1,2}} \underline{a_{2,3}b_{1,2}} \underline{a_{1,3}a_{2,3}} \simeq a_{1,3} \underline{a_{1,2}a_{1,2}} b_{1,2} a_{2,3} a_{1,3} b_{1,2} a_{2,3} \\
&\simeq \underline{a_{1,3}b_{1,2}} \underline{a_{2,3}a_{1,3}} \underline{b_{1,2}a_{2,3}} \simeq a_{2,3}a_{2,3} \simeq \emptyset.
\end{aligned}$$

5. Let $a \in \alpha$ be a letter such that $\tau(a) \neq a$. Set $b = \tau(a)$. Then the words $aabab$, $babaa$, and $baaab$ are pairwise homotopic and homotopic to the nanoword $w = AA'AA'$ with $|A| = |A'| = a$:

$$(aabab)^d = a_{1,2}a_{1,3}a_{1,2} \underline{a_{2,3}b_{1,2}} \underline{a_{1,3}a_{2,3}b_{1,2}} \simeq a_{1,2}a_{1,3}a_{1,2}a_{1,3} \approx w,$$

$$(babaa)^d = \underline{b_{1,2}a_{1,2}} \underline{a_{1,3}b_{1,2}a_{1,2}} \underline{a_{2,3}a_{1,3}a_{2,3}} \simeq a_{1,3}a_{2,3}a_{1,3}a_{2,3} \approx w.$$

$$(baaab)^d = b_{1,2} \underline{a_{1,2}a_{1,3}} \underline{a_{1,2}a_{2,3}} \underline{a_{1,3}a_{2,3}b_{1,2}} \simeq \underline{b_{1,2}a_{1,3}} \underline{a_{1,2}a_{2,3}a_{1,2}a_{2,3}} \underline{a_{1,3}b_{1,2}} \simeq w.$$

The word $aabab$ is not symmetric but is homotopically symmetric since $(aabab)^- = babaa \simeq aabab$.

3.5. Remarks. 1. In analogy with the theory of framed knots in topology, one can define a *framed* homotopy move on nanowords $(\mathcal{A}, xAAyBBz) \mapsto (\mathcal{A}, xyz)$ which applies when $|A| = \tau(|B|)$. This move together with homotopy moves of types 2 and 3 generates an equivalence relation on nanowords called *framed homotopy*. It is stronger than homotopy. We shall not study framed homotopy here.

2. One can generalize the second homotopy move $(\mathcal{A}, xAB yBAz) \mapsto (\mathcal{A} - \{A, B\}, xyz)$ allowing it whenever the pair $(|A|, |B|)$ lies in a fixed subset of α^2 . We study here only the case where this subset is the graph of τ .

3. A *homotopy automorphism* of an étale word w over α is a τ -equivariant permutation $f : \alpha \rightarrow \alpha$ such that $f_{\#}(w) \simeq w$. The classification theorems below allow to compute the group of homotopy automorphisms for words of length ≤ 5 and nanowords of length ≤ 6 .

4. It is possible that all étale words over a 1-element alphabet are contractible; at least I do not know obstructions to this.

4. HOMOMORPHISM γ

We introduce an elementary homotopy invariant of nanowords over α . The idea is to associate with a nanoword an element of a group generated by elements of α .

4.1. Group Π and homomorphism γ . Recall that an *orbit* of the involution $\tau : \alpha \rightarrow \alpha$ is a subset of α consisting either of one element preserved by τ or of two elements permuted by τ ; in the latter case the orbit is *free*. Let Π be the group with generators $\{z_a\}_{a \in \alpha}$ and defining relations $z_a z_{\tau(a)} = 1$ for $a \in \alpha$. It is clear that Π is a free product of k infinite cyclic groups and l cyclic groups of order 2 where k is the number of free orbits of τ and l is the number of fixed points of τ . The formula $z_a \mapsto z_{\tau(a)}$ defines an involutive automorphism of Π denoted τ_* .

Consider a nanoword $(\mathcal{A}, w : \hat{n} \rightarrow \mathcal{A})$ over α . For $i = 1, \dots, n$, set $\gamma_i = z_{|w(i)|}$ if i numerates the first entry of $w(i)$ in w , that is if $w(i) \neq w(j)$ for $j < i$. Otherwise, set $\gamma_i = z_{\tau(|w(i)|)} = (z_{|w(i)|})^{-1}$. Set $\gamma(w) = \gamma_1 \cdots \gamma_n \in \Pi$. For example, if $w = ABAB$ with $|A| = a \in \alpha, |B| = b \in \alpha$, then $\gamma(w) = z_a z_b z_a^{-1} z_b^{-1}$. It is easy to check that $\gamma(w)$ is invariant under homotopy moves on w . This provides an efficient and easily computable homotopy invariant of nanowords. The mapping $w \mapsto \gamma(w)$ defines a monoid homomorphism $\gamma : \mathcal{N}_\bullet(\alpha) \rightarrow \Pi$ such that $\gamma(\overline{w}) = \tau_*(\gamma(w))$ and $\gamma(w^-) = (\gamma(w))^{-1}$ for any w .

Lemma 4.1.1. $\gamma(\mathcal{N}_\bullet(\alpha)) = [\Pi, \Pi]$.

Proof. It is obvious that the composition of γ with the projection $\Pi \rightarrow \Pi/[\Pi, \Pi]$ is a trivial homomorphism. Hence $\gamma(\mathcal{N}_\bullet(\alpha)) \subset [\Pi, \Pi]$. To prove the opposite inclusion, pick a set $\alpha_0 \subset \alpha$ meeting each orbit of τ in one element. The group Π can be presented by the generators $\{z_a\}_{a \in \alpha_0}$ and defining relations $z_a z_a = 1$ for all $a \in \alpha_0$ such that $\tau(a) = a$. Each element of $[\Pi, \Pi]$ can be presented by a word W in the alphabet $\{z_a^{\pm 1}\}_{a \in \alpha_0}$ in which every z_a appears with total power 0. Partition the entries of $z_a^{\pm 1}$ in W into pairs (z_a, z_a^{-1}) or (z_a^{-1}, z_a) in an arbitrary way and let s_a be the resulting set of pairs. For a pair $A \in s_a$, set $|A| = a$ if the first entry of this pair is z_a and set $|A| = \tau(a)$ if the first entry of this pair is z_a^{-1} . This makes the set $\mathcal{A} = \Pi_a s_a$ into an α -alphabet. Replacing in W each entry $z_a^{\pm 1}$ with the only $A \in s_a$ containing this entry we obtain a nanoword $(\mathcal{A}, w)$ such that $\gamma(w) = W$. For example if $W = z_a^2 z_b^{-1} z_a^{-1} z_b z_a^{-1}$, then $(\mathcal{A} = \{A_1, A_2, B\}, w = A_1 A_2 B A_1 B A_2)$ with $|A_1| = |A_2| = a, |B| = \tau(b)$ is a nanoword such that $\gamma(w) = W$. $\square$

The commutator group $[\Pi, \Pi]$ is a free group for any τ . Its rank is infinite if (in the notation above) $k \geq 2$ or $k = 1$ and $l \geq 1$. If $k = 1$ and $l = 0$, then $[\Pi, \Pi] = 1$. If $k = 0$, then $[\Pi, \Pi]$ is a free group of rank $2^{l-1}(l-2) + 1$. One way to see it goes by realizing Π as the fundamental group of the connected sum X of l copies of RP^3 and observing that the maximal abelian covering of X has the same fundamental group as a connected graph with $2^{l-1}l$ vertices and $2^l(l-1)$ edges.

If τ has at least two orbits, then Lemma 4.1.1 implies that $\mathcal{N}_\bullet(\alpha)$ is an infinite monoid. If τ has at least three orbits, then Lemma 4.1.1 implies that $\mathcal{N}_\bullet(\alpha)$ is non-abelian.

4.2. Homomorphism γ' . It is sometimes sufficient to consider a weaker invariant $\gamma' : \mathcal{N}_\bullet(\alpha) \rightarrow \Pi'$ where Π' is obtained from Π by adding relations $z_a z_a = 1$ for all a and γ' is the composition of γ with the projection $\Pi \rightarrow \Pi'$. To give an example, pick an integer $m \geq 1$ and elements $a, b \in \alpha$ lying in different orbits of τ . Then $z_a z_b \in \Pi'$ is an element of infinite order and $(z_a z_b)^{-1} = z_b z_a$ in Π' . Consider the nanoword $w_m = A_1 B_1 A_2 B_2 \cdots A_m B_m A_1 B_1 A_2 B_2 \cdots A_m B_m$ with $m \geq 1$ and $|A_i| = a, |B_i| = b$ for all i . It is clear that $\gamma'(w_m) = (z_a z_b)^{2m}$. Therefore the nanowords $\{w_m\}_m$ are non-contractible and mutually non-homotopic. The opposite nanowords $w_m^- = B_m A_m \cdots B_1 A_1 B_m A_m \cdots B_1 A_1$ are also mutually non-homotopic and homotopically distinct from $\{w_m\}_m$. Indeed $\gamma'(w_m^-) = (\gamma'(w_m))^{-1} = (z_b z_a)^{2m} = (z_a z_b)^{-2m}$ for any $m \geq 1$.

We use γ' to associate with any nanoword w a function $\mu_w : \hat{\alpha} \times \hat{\alpha} \rightarrow \mathbb{Z}$ where $\hat{\alpha} = \alpha/\tau$ is the set of orbits of τ . The orbit $\{a, \tau(a)\}$ of $a \in \alpha$ will be denoted $\hat{a}$. Pick $a, b \in \alpha$. If $\hat{a} = \hat{b}$, then $\mu_w(\hat{a}, \hat{b}) = 0$. Suppose that a and b lie in different orbits. Consider the group $\Pi_0 = (x, y \mid x^2 = y^2 = 1)$ and the group homomorphism $\rho : \Pi' \rightarrow \Pi_0$ such that $\rho(z_a) = \rho(z_{\tau(a)}) = x$, $\rho(z_b) = \rho(z_{\tau(b)}) = y$, and $\rho(z_c) = 1$ for all other $c \in \alpha$. Observe that $[\Pi_0, \Pi_0]$ is an infinite cyclic group generated by $xyxy = (yxyx)^{-1}$. Since $\rho\gamma'(w) \in [\Pi_0, \Pi_0]$, there is a unique $m \in \mathbb{Z}$ such that $\rho\gamma'(w) = (xyxy)^m$. Set $\mu_w(\hat{a}, \hat{b}) = m$. It follows from the definitions that the mapping $\mu_w : \hat{\alpha} \times \hat{\alpha} \rightarrow \mathbb{Z}$ is skew-symmetric in the sense that $\mu_w(\hat{a}, \hat{b}) = -\mu_w(\hat{b}, \hat{a})$ for all $a, b \in \alpha$. Since $\gamma'(w)$ is a homotopy invariant of w , so is μ_w . Note that $\mu_{w_1 w_2} = \mu_{w_1} + \mu_{w_2}$ for any nanowords w_1, w_2 . For $w = ABAB$ with $|A| = a \in \alpha, |B| = b \in \alpha$ lying in different orbits of τ , the function μ_w takes values 1 and -1 on the pairs $\hat{a}, \hat{b}$ and $\hat{b}, \hat{a}$, respectively, and the value 0 on all other pairs. This implies that every skew-symmetric mapping $\hat{\alpha} \times \hat{\alpha} \rightarrow \mathbb{Z}$ is the μ -function of a nanoword over α .

4.3. Homomorphism $\tilde{\gamma}$. A more careful approach yields the following refined version of γ . Let $\tilde{\Pi}$ be the group with generators $\{\tilde{z}_a\}_{a \in \alpha}$ and defining relations $\tilde{z}_a \tilde{z}_{\tau(a)} \tilde{z}_b = \tilde{z}_b \tilde{z}_a \tilde{z}_{\tau(a)}$ for all $a, b \in \alpha$. The formula $\tilde{z}_a \mapsto z_a$ defines a projection $\tilde{\Pi} \rightarrow \Pi$. Replacing z_a with $\tilde{z}_a$ in the definition of γ , we obtain a lift of γ to a monoid homomorphism $\tilde{\gamma} : \mathcal{N}_\bullet(\alpha) \rightarrow \tilde{\Pi}$. Clearly, $\tilde{\gamma}(\mathcal{N}_\bullet(\alpha)) \subset [\tilde{\Pi}, \tilde{\Pi}]$.

Despite many nice features, the homomorphisms γ , γ' , and $\tilde{\gamma}$ often fail to distinguish homotopy classes of nanowords. For example, as we shall see in Sect. 8.4, the nanoword $w = ABAB$ with $|A| = |B| \neq \tau(|A|)$ is non-contractible but obviously $\tilde{\gamma}(w) = 1$.

5. COVERINGS OF NANOWORDS

We define for each nanoword a family of nanowords called its coverings.

5.1. Group π and interlacement of letters. Let $\pi = \pi(\alpha, \tau)$ be the multiplicative abelian group with generators $\{a\}_{a \in \alpha}$ and defining relations $a\tau(a) = 1$ for all $a \in \alpha$. Clearly, $\pi = \Pi/[\Pi, \Pi]$ where Π is the group considered in Sect. 4.1. The group π is a direct product of copies of $\mathbb{Z}/2\mathbb{Z}$ numerated by $\{a \in \alpha \mid \tau(a) = a\}$ and copies of $\mathbb{Z}$ numerated by free orbits of τ .

Consider a nanoword $(\mathcal{A}, w)$ over α . We say that two letters $A, B \in \mathcal{A}$ are *w-interlaced* if

$$w = \dots A \dots B \dots A \dots B \dots \quad \text{or} \quad w = \dots B \dots A \dots B \dots A \dots.$$

Set in the first case $n_w(A, B) = 1 \in \mathbb{Z}$ and in the second case $n_w(A, B) = -1 \in \mathbb{Z}$. For all other pairs of letters $A, B \in \mathcal{A}$, we set $n_w(A, B) = 0 \in \mathbb{Z}$ and say that A, B are *w-unlaced*. Note that $n_w(A, B) = -n_w(B, A)$ and $n_w(A, A) = 0$. For any $A \in \mathcal{A}$, set

$$[A]_w = \prod_{B \in \mathcal{A}} |B|^{n_w(A, B)} \in \pi.$$

5.2. Coverings. By an α -family of subgroups of π , we mean a family $H = \{H_a \subset \pi\}_{a \in \alpha}$ where H_a is a subgroup of π such that $H_a = H_{\tau(a)}$ for all a . For a nanoword $(\mathcal{A}, w)$ over α , consider the nanoword $(\mathcal{A}^H, w^H)$ over α obtained from $(\mathcal{A}, w)$ by deleting from both $\mathcal{A}$ and w all letters A such that $[A]_w \notin H_{|A|}$. It is called the *H-covering* of $(\mathcal{A}, w)$.

Lemma 5.2.1. *If two nanowords are homotopic, then their H-coverings are homotopic.*

Proof. It is clear that H -coverings of isomorphic nanowords are isomorphic. It remains to prove that if a nanoword v is obtained from a nanoword w by a homotopy move then v^H is obtained from w^H by a homotopy move. Consider the move $w = (\mathcal{A}, xAAy) \mapsto (\mathcal{A} - \{A\}, xy) = v$. Then $n_w(A, B) = n_w(B, A) = 0$ for all $B \in \mathcal{A}$. Hence $[B]_w = [B]_v$ for any $B \in \mathcal{A} - \{A\}$ and $[A]_w = 1 \in \pi$. Therefore A survives in w^H and the nanoword v^H is obtained from w^H by deleting A , i.e., by the first homotopy move.

Consider the move $w = (\mathcal{A}, xAB yBAz) \mapsto (\mathcal{A} - \{A, B\}, xyz) = v$ where $|A| = a \in \pi$ and $|B| = \tau(a)$. It follows from the definitions that $n_w(A, B) = 0$ and $n_w(A, C) = n_w(B, C)$ for all $C \in \mathcal{A} - \{A, B\}$. Therefore $[A]_w = [B]_w$. Observe also that for $C \in \mathcal{A} - \{A, B\}$,

$$[C]_w [C]_v^{-1} = |A|^{n_w(C, A)} |B|^{n_w(C, B)} = (a\tau(a))^{n_w(C, A)} = 1.$$

Hence $[C]_w = [C]_v$. Therefore in the case $[A]_w \notin H_a$, we have $w^H = v^H$ and in the case $[A]_w \in H_a$, the nanoword v^H is obtained from w^H by the second homotopy move.

Consider the move $w = (\mathcal{A}, xAB yAC zBCt) \mapsto (\mathcal{A}, xBA yCA zCBt) = v$. Set $a = |A| = |B| = |C| \in \alpha$. It is clear that for $D \in \mathcal{A} - \{A, B, C\}$ and $E \in \mathcal{A}$, we have $n_w(D, E) = n_v(D, E)$. Thus $[D]_w = [D]_v$. Therefore D either disappears in both w^H, v^H or survives in both w^H, v^H . For $D \in \{A, B, C\}$, the sum $n_w(D, A) + n_w(D, B) + n_w(D, C)$ is preserved under the move. The product

$$|A|^{n_w(D, A)} |B|^{n_w(D, B)} |C|^{n_w(D, C)} = a^{n_w(D, A) + n_w(D, B) + n_w(D, C)}$$

is also preserved. This implies that $[D]_w = [D]_v$. We claim that $[A]_w[C]_w = [B]_w$. To see this, set $x * y = \prod_E [E]_w \in \pi$ where E runs over the letters which appear both in x and in y . The expressions $y * z, y * t$, etc. are defined similarly. It follows from the definitions that

$$\begin{aligned} [A]_w &= a^{n_w(A,A)+n_w(A,B)+n_w(A,C)}(y * z)(y * t)(x * y)^{-1} = a(y * z)(y * t)(x * y)^{-1}, \\ [B]_w &= a^{n_w(B,A)+n_w(B,B)+n_w(B,C)}(y * t)(z * t)(x * y)^{-1}(x * z)^{-1} \\ &= (y * t)(z * t)(x * y)^{-1}(x * z)^{-1}, \end{aligned}$$

$$[C]_w = a^{n_w(C,A)+n_w(C,B)+n_w(C,C)}(z * t)(x * z)^{-1}(y * z)^{-1} = a^{-1}(z * t)(x * z)^{-1}(y * z)^{-1}.$$

These computations imply that $[A]_w[C]_w = [B]_w$. There are three possibilities: the elements $[A]_w, [B]_w, [C]_w \in \pi$ belong to H_a ; exactly one of them belongs to H_a ; neither of them belongs to H_a . In the first case v^H is obtained from w^H by the third homotopy move. In the second and third cases $v^H = w^H$. $\square$

Note that $(w^-)^H = (w^H)^-, \overline{w^H} = \overline{w^H}, (ww')^H = w^H(w')^H$ for any nanowords w, w' .

Any τ -invariant subgroup H of π determines an α -family of subgroups of π by $H_a = H$ for all $a \in \alpha$. This family is denoted by the same symbol H .

The following example shows that homotopy invariants of the coverings can be efficiently used to distinguish homotopy classes of nanowords.

5.3. Example. Set $\alpha = \{a, b\}$ with $\tau = \text{id}$. Consider the nanoword $w = A_1 B_1 B_2 A_2 A_1 A_3 B_1 A_3 A_2 B_2$ with $|A_i| = a$ and $|B_j| = b$ for all i, j . We have $\pi = \pi(\alpha, \tau) = (a, b | a^2 = b^2 = 1, ab = ba)$, $[A_1]_w = a \in \pi, [A_3]_w = b \in \pi$ and $[A_2]_w = [B_1]_w = [B_2]_w = ab \in \pi$. Let $H = \{1, ab\} \subset \pi$. Then $w^H = B_1 B_2 A_2 B_1 A_2 B_2$. Clearly, $\gamma(w^H) = z_a z_b z_a z_b \neq 1$ and $\mu_{w^H}(a, b) = 1$. Thus w is non-contractible. All the invariants of w defined in Sect. 4 are trivial since $\tilde{\gamma}(w) = 1$.

6. SELF-LINKING

Starting from the notion of linking (or interlacement) of letters in a word, we construct a family of polynomial invariants of nanowords.

6.1. Self-linking class. Let $\pi = \pi(\alpha, \tau)$ be the group defined in Sect. 5.1. Recall that given a nanoword $(\mathcal{A}, w)$ over α , we associate with every $A \in \mathcal{A}$ an element $[A]_w$ of π , see Sect. 5.1. Set $\mathcal{A}_w = \{A \in \mathcal{A} | [A]_w \neq 1\}$. For a letter $a \in \alpha$, define its *self-linking class* by

$$[a]_w = \sum_{A \in \mathcal{A}_w, |A|=a} [A]_w \in \mathbb{Z}\pi$$

where $\mathbb{Z}\pi$ is the group ring of π with integer coefficients. By definition, the neutral element $1 \in \pi$ appears in this expression with coefficient 0. The key property of the self-linking class is provided by the following theorem.

Theorem 6.1.1. *If $\tau(a) = a$, then $[a]_w(\text{mod } 2) \in (\mathbb{Z}/2\mathbb{Z})\pi$ is a homotopy invariant of w . If $\tau(a) \neq a$, then $[a]_w - [\tau(a)]_w \in \mathbb{Z}\pi$ is a homotopy invariant of w .*

Proof. It is obvious that isomorphic nanowords give rise to the same self-linking class for all $a \in \alpha$. Consider the first homotopy move $w = (\mathcal{A}, xAAy) \mapsto (\mathcal{A} - \{A\}, xy) = v$. As was shown in the proof of Lemma 5.2.1, $[B]_w = [B]_v$ for any $B \in \mathcal{A} - \{A\}$ and $[A]_w = 1$. Therefore $[a]_w = [a]_v$ for any $a \in \alpha$.

Consider the move $w = (\mathcal{A}, xAB yBAz) \mapsto (\mathcal{A} - \{A, B\}, xyz) = v$ where $|A| = \tau(|B|)$. By the proof of Lemma 5.2.1, $[A]_w = [B]_w$ and $[C]_w = [C]_v$ for $C \in \mathcal{A} - \{A, B\}$. This implies that if $a \in \alpha - \{|A|, |B|\}$, then $[a]_w = [a]_v$. If $a = |A| = |B|$, then A, B contribute $[A]_w + [B]_w = 2[A]_w$ to $[a]_w$ so that $[a]_w = [a]_v(\text{mod } 2)$. If $a = |A| \neq \tau(a) = |B|$ then A, B contribute $[A]_w - [B]_w = 0$ to $[a]_w - [\tau(a)]_w$ so that $[a]_w - [\tau(a)]_w = [a]_v - [\tau(a)]_v$.

Consider the move $w = (\mathcal{A}, xAB yACzBCt) \mapsto (\mathcal{A}, xBA yCAzCBt) = v$. By the proof of Lemma 5.2.1, $[D]_w = [D]_v$ for any $D \in \mathcal{A}$. Hence $[a]_w = [a]_v$ for all $a \in \alpha$. $\square$

6.2. Self-linking function. Theorem 6.1.1 suggests the following definitions. For $a \in \alpha$, set $R_a = \mathbb{Z}/2\mathbb{Z}$ if $\tau(a) = a$ and $\mathbb{Z}_a = \mathbb{Z}$ if $\tau(a) \neq a$. Consider the system of commutative rings $\{R_a\pi\}_{a \in \alpha}$. By a *section* of this system we mean a mapping which assigns to every $a \in \alpha$ an element of $R_a\pi$. Each nanoword w determines such a section u^w by $u^w(a) = [a]_w(\text{mod } 2) \in R_a\pi$ if $\tau(a) = a$ and $u^w(a) = [a]_w - [\tau(a)]_w \in R_a\pi$ if $\tau(a) \neq a$. By Theorem 6.1.1, u^w , also denoted $u(w)$, is a homotopy invariant of w . It is called the *self-linking function* of w . If w is contractible, then $u^w = 0$.

It is clear that $u^w(\tau(a)) = -u^w(a)$ for all a . Therefore the function u^w is entirely determined by its restriction to any set $\alpha_0 \subset \alpha$ meeting every orbit of τ in exactly one element. We call such α_0 an *orientation* of τ . Given α_0 and $a \in \alpha_0$, we can view $u^w(a)$ as a Laurent polynomial as follows. Any element of π expands as $\prod_{b \in \alpha_0} b^{m_b}$ with $m_b \in \mathbb{Z}$. The integer m_b is uniquely determined if $\tau(b) \neq b$ and is determined modulo 2 if $\tau(b) = b$. The elements of $R_a\pi$ (and in particular $u^w(a)$) are just Laurent polynomials with coefficients in R_a in the commuting variables $b \in \alpha_0$ subject to $b^2 = 1$ for $\tau(b) = b$.

For a nanoword w obtained as the product of nanowords w_1 and w_2 , we have $u^w(a) = u^{w_1}(a) + u^{w_2}(a)$ for all $a \in \alpha$. This follows from the definitions and the fact that letters of w_1 and w_2 are unlaced in w . Thus the formula $w \rightarrow u^w$ defines a monoid homomorphism from $\mathcal{N}_\bullet(\alpha)$ to the additive group of sections of the system $\{R_a\pi\}_{a \in \alpha}$. We describe the image of this homomorphism in the next subsection.

To compute the self-linking function of the opposite nanoword w^- , observe that $[a]_{w^-} = \overline{[a]_w}$ for any $a \in \alpha$ where the overbar denotes the involution in $R_a\pi$ sending elements of π to their inverses. Therefore $u(w^-) = \overline{u(w)}$. For the inverse nanoword $\overline{w}$, we have the identity $[a]_{\overline{w}} = \overline{[\tau(a)]_w}$. Therefore $u(\overline{w}) = -\overline{u(w)}$ and $u(\overline{w}^-) = -u(w)$. In other words $u(w\overline{w}^-) = 0$ for all w . If w is homotopically skew-symmetric, then $u^w(a) = 0$ for all $a \in \alpha$ such that $\tau(a) \neq a$.

The self-linking function can be used to estimate $\|w\|$ from below. If w is homotopic to a nanoword of length $2m$, then for any $a \in \alpha$, the element $u^w(a) \in R_a\pi$ is an algebraic sum of monomials in the generators $\{b\}_{b \in \alpha} \subset \pi$ of degree $\leq m - 1$.

6.3. Characterization. Consider the following characterization problem: given an orientation $\alpha_0 \subset \alpha$, when a system $u = \{u(a) \in R_a\pi\}_{a \in \alpha_0}$ is the self-linking function of a nanoword? We call such a system a *section* of $\{R_a\pi\}_{a \in \alpha_0}$. Denote by δ_a the additive homomorphism $R_a\pi \rightarrow R_a$ sending $1 \in \pi$ to $1 \in R_a$ and sending all other elements of π to 0. For $a, b \in \alpha_0$, set $R_{a,b} = \mathbb{Z}$ if $\tau(a) \neq a$ and $\tau(b) \neq b$ and $R_{a,b} = \mathbb{Z}/2\mathbb{Z}$ otherwise. (Clearly $R_{a,b} = R_a \otimes R_b$ and $R_{a,a} = R_a$). Let ∂_b be the additive homomorphism $R_a\pi \rightarrow R_{a,b}$ sending a monomial $\prod_{b \in \alpha_0} b^{m_b}$ to m_b . We say that a section u of $\{R_a\pi\}_{a \in \alpha_0}$ is *skew-symmetric* if $\delta_a(u(a)) = \partial_a(u(a)) = 0$ for any $a \in \alpha_0$ and $\partial_a(u(b)) + \partial_b(u(a)) = 0$ for any $a, b \in \alpha_0$.

Theorem 6.3.1. *A section of $\{R_a\pi\}_{a \in \alpha_0}$ is the self-linking function of a nanoword if and only if it is skew-symmetric.*

Proof. Pick a nanoword w and set $u = u^w$. The equality $\delta_a(u(a)) = 0$ follows from the definitions. Let us prove the other identities. For $a, b \in \alpha_0$, set

$$a \circ b = \sum_{A, B \in \mathcal{A}, |A|=a, |B|=b} n_w(A, B) \in R_{a,b}.$$

The identities $n_w(A, B) + n_w(B, A) = 0$ and $n_w(A, A) = 0$ imply that $a \circ b + b \circ a = a \circ a = 0$.

Suppose first that $\tau(a) = a, \tau(b) = b$. Observe that for any $A \in \mathcal{A} - \mathcal{A}_w$ we have $\partial_b([A]_w) = 0$. Therefore

$$\begin{aligned} \partial_b(u(a)) &= \partial_b([a]_w) = \sum_{A \in \mathcal{A}_w, |A|=a} \partial_b([A]_w) = \sum_{A \in \mathcal{A}, |A|=a} \partial_b([A]_w) \\ &= \sum_{A \in \mathcal{A}, |A|=a} \partial_b\left(\prod_{B \in \mathcal{A}} |B|^{n_w(A, B)}\right) = \sum_{A, B \in \mathcal{A}, |A|=a, |B|=b} n_w(A, B) = a \circ b. \end{aligned}$$

This implies that $\partial_a(u(a)) = 0 = \partial_a(u(b)) + \partial_b(u(a))$.

Suppose that $\tau(a) = a, \tau(b) \neq b$. Then as above

$$\begin{aligned} \partial_b(u(a)) &= \partial_b([a]_w) = \sum_{A \in \mathcal{A}, |A|=a} \partial_b\left(\prod_{B \in \mathcal{A}} |B|^{n_w(A, B)}\right) \\ &= \sum_{A, B \in \mathcal{A}, |A|=a, |B|=b} n_w(A, B) - \sum_{A, B \in \mathcal{A}, |A|=a, |B|=\tau(b)} n_w(A, B) = a \circ b - a \circ \tau(b), \\ \partial_a(u(b)) &= \partial_a([b]_w) = \partial_a([b]_w - [\tau(b)]_w) \\ &= \sum_{B \in \mathcal{A}, |B|=b} \partial_a([B]_w) - \sum_{B \in \mathcal{A}, |B|=\tau(b)} \partial_a([B]_w) = b \circ a - \tau(b) \circ a. \end{aligned}$$

Hence

$$\partial_a(u(b)) + \partial_b(u(a)) = b \circ a - \tau(b) \circ a + a \circ b - a \circ \tau(b) = 0.$$

In the case $\tau(a) \neq a, \tau(b) \neq b$, similar computations give

$$\partial_b(u(a)) = a \circ b - a \circ \tau(b) - \tau(a) \circ b + \tau(a) \circ \tau(b)$$

and the equalities $\partial_a(u(a)) = 0 = \partial_a(u(b)) + \partial_b(u(a))$ follow from the properties of $\circ$.

We now prove the sufficiency of the conditions stated in the theorem. For $a \in \alpha$, we call an element of $R_a\pi$ *linear* if it is a linear combination of elements of $\alpha_0 \subset \pi$. (Elements of $\alpha - \alpha_0 \subset \pi$ and $1 \in R_a\pi$ are not linear). A section u of $\{R_a\pi\}_{a \in \alpha_0}$ is *linear* if its values on all elements of α_0 are linear. A section u of $\{R_a\pi\}_{a \in \alpha_0}$ is *realizable* if it is the restriction to α_0 of the self-linking function of a nanoword. Note that if u is realizable then so is $-u$. By the additivity of the self-linking function, the sum of realizable systems is realizable.

The idea of the proof is to take any skew-symmetric system u and to deduce from it consecutively certain “model” realizable sections so that at the end we obtain 0. Since 0 is realized by an empty nanoword, u is realizable. A model section is defined as follows. Pick $a \in \alpha_0, b_1, \dots, b_n \in \alpha$ (possibly with repetitions) where $n \geq 1$. Consider the nanoword $w = B_1 B_2 \cdots B_n A B_n \cdots B_2 B_1 A$ with $|A| = a$ and $|B_i| = \tau(b_i)$ for all i . It is clear that $[A]_w = \prod_i b_i$ and $[B_i]_w = a$ for all i . It follows from the definitions that $u^w(a) = \prod_i b_i$ modulo linear terms (corresponding to i such that $b_i = a$ or $b_i = \tau(a)$). The values of u^w on all elements of $\alpha_0 - \{a\}$ are linear. Adding or subtracting such model sections u^w from the original section u and using that $\delta_a(u(a)) = 0$ we can obtain a linear skew-symmetric section, again denoted u . For any $a \in \alpha_0$, we can uniquely expand $u(a) = \sum_{b \in \alpha_0} r(a, b)b$ where $r(a, b) \in R_a$. Pick $a \in \alpha_0$ such that $\tau(a) = a$. For $b \in \alpha_0 - \{a\}$, consider the nanoword $w = ABAB$ with $|A| = a$ and $|B| = b$. It is clear that $[A]_w = b$ and $[B]_w = a^{-1} = a \in \pi$. Therefore $u^w(a) = b, u^w(b) = a$, and all other values of u^w are 0. Adding $\pm u^w$ to u , we change $r(a, b)$ by ± 1 keeping all $r(a, b')$ for $b' \neq b$. Proceeding by induction, we transform u into a linear skew-symmetric section, still denoted u , such that $r(a, b) = 0$ for all $b \in \alpha - \{a\}$. By skew-symmetry, $r(b, a) = r(a, b) = 0$ and $r(a, a) = \partial_a(a) = 0 \in \mathbb{Z}/2\mathbb{Z}$. Thus $u(a) = 0$ and a does not show up in the expansions of $u(b)$ for all $b \in \alpha_0$. Inductively, we can ensure that this holds for all $a \in \alpha_0$ such that $\tau(a) = a$. Pick now $a \in \alpha_0$ with $\tau(a) \neq a$. Pick $b \in \alpha_0 - \{a\}$ with $\tau(b) \neq b$. Consider the nanoword $w = ABAB$ with $|A| = \tau(a)$ and $|B| = b$. It is clear that $u^w(a) = -[A]_w = -b, u^w(b) = [B]_w = (\tau(a))^{-1} = a$. Adding the linear section $\pm u^w$ to u , we change $u(a)$ by $\pm b$ keeping all other values of u except $u(b)$. Proceeding in this way we can ensure that $u(a) = r(a, a)a$. By skew-symmetry, $r(a, a) = \partial_a(a) = 0 \in \mathbb{Z}$. Then $u(a) = 0$ and again by skew-symmetry, $r(b, a) = 0$ for all $b \in \alpha_0 - \{a\}$. Proceeding further in this way we eventually transform u into the zero section as required. $\square$

6.4. Examples. 1. Pick $a \in \alpha$ with $\tau(a) \neq a$ and consider the nanoword $w = ABAB$ with $|A| = |B| = a$. Clearly, $[A]_w = a, [B]_w = a^{-1}$ and the self-linking function u^w takes the value $a + a^{-1} \in R_a\pi = \mathbb{Z}\pi$ on a , the value $-a - a^{-1}$ on $\tau(a)$, and the value 0 on all other elements of α . Since $u^w \neq 0$, the word w is not

contractible. Since $u^w \neq -u^w$, the word w is not homotopic to its inverse. For $n \geq 2$, the polynomial $u^w(a)$ is not divisible by n and therefore w is not homotopic to the n -th power of a nanoword. Note that $\overline{u(w)} = u(w)$ which is compatible with the fact that w is isomorphic to its opposite.

2. Fix $a, b \in \alpha$ belonging to different orbits of τ . For finite sequences $\varepsilon = (\varepsilon_i)_{i=1}^m, \mu = (\mu_j)_{j=1}^n$ consisting of zeros and ones we define a nanoword

$$w = w(\varepsilon, \mu) = A_1 A_2 \cdots A_m B_1 B_2 \cdots B_n A_m \cdots A_2 A_1 B_n \cdots B_2 B_1$$

over α with $|A_i| = \tau^{\varepsilon_i}(a)$ and $|B_j| = \tau^{\mu_j}(b)$ for all i, j . We give a homotopy classification of these nanowords using the self-linking function.

Case $\tau(a) = a, \tau(b) = b$. Observe that $u^w(a) = 0$ if mn is even and $u^w(a) = b$ if mn is odd. Hence $mn \pmod{2}$ is a homotopy invariant of $w = w(\varepsilon, \mu)$. Applying the first two homotopy moves it is easy to see that $mn \pmod{2}$ is a full homotopy invariant of $w(\varepsilon, \mu)$: any two nanowords of this type with the same $mn \pmod{2}$ are homotopic.

Case $\tau(a) \neq a, \tau(b) = b$. By the second homotopy move, if the length n of μ is even then $w = w(\varepsilon, \mu)$ is contractible. Suppose that n is odd. Set

$$k = \text{card} \{i = 1, \dots, m \mid \varepsilon_i = 0\} - \text{card} \{i = 1, \dots, m \mid \varepsilon_i = 1\} \in \mathbb{Z}.$$

A direct computation gives $u^w(a) = kb$ so that k is a homotopy invariant. It is easy to see that k is a full homotopy invariant: two nanowords $w(\varepsilon, \mu), w(\varepsilon', \mu')$ with μ, μ' of odd length are homotopic if and only if they give rise to the same k . In particular, such a nanoword is contractible iff $k = 0$.

Case $\tau(a) = a, \tau(b) \neq b$. This case is similar to the previous one. If the length m of ε is even then $w(\varepsilon, \mu)$ is contractible. For odd m , the classifying invariant is the integer

$$l = \text{card} \{j = 1, \dots, m \mid \mu_j = 0\} - \text{card} \{j = 1, \dots, m \mid \mu_j = 1\}.$$

Case $\tau(a) \neq a, \tau(b) \neq b$. Let k, l be the integers determined by ε, μ as above. We have $u^w(a) = kb^l$ if $l \neq 0$ and $u^w(a) = 0$ if $l = 0$. Similarly, $u^w(b) = la^{-k}$ if $k \neq 0$ and $u^w(b) = 0$ if $k = 0$. Therefore $w(\varepsilon, \mu)$ is contractible iff $k = 0$ or $l = 0$. For the nanowords $w(\varepsilon, \mu)$ with $kl \neq 0$, the pair (k, l) is a full homotopy invariant: two such nanowords are homotopic if and only if they give rise to the same k, l .

3. The following example illustrates the homotopy invariants derived from the self-linking function via coverings. Let $\alpha = \{a, b\}$ with involution τ permuting a and b . The corresponding group π is an infinite cyclic group generated by a and $b = a^{-1}$. The self-linking function u^w of any nanoword w over α is determined by the Laurent polynomial $u^w(a) \in \mathbb{Z}\pi = \mathbb{Z}[a^{\pm 1}]$. For integers $m, n \geq 1$, consider the nanoword

$$w = w_{m,n} = A_1 A_2 \cdots A_m B_1 B_2 \cdots B_n A_m \cdots A_2 A_1 B_n \cdots B_2 B_1.$$

with $|A_i| = a$ and $|B_j| = b$ for all i, j . It is clear that $[A_i]_w = b^n$ and $[B_j]_w = a^{-m}$ for all i, j . Therefore

$$u^w(a) = [a]_w - [\tau(a)]_w = [a]_w - [b]_w = mb^n - na^{-m} = ma^{-n} - na^{-m}.$$

Pick integers $p, q, s, t \geq 1$ and consider the nanoword v over α of length $12(p + q + s + t)$ obtained as the product (in an arbitrary order) of the nanowords

$$w_{p,s}, w_{p,t}, w_{q,s}, w_{q,t}, w_{s+t,p}, w_{s+t,q}, w_{s,p+q}, w_{t,p+q}, w_{p+q,s+t}.$$

The additivity of the self-linking function easily implies that $u(v) = 0$. Let r be the greatest common divisor of p and s . Let H be the subgroup of π generated by a^r . Suppose that $r \geq 2$ and q, t are prime to r . It is easy to check that $v^H \simeq w_{p,s}$ and therefore $u(v^H) = pa^{-s} - sa^{-p} \neq 0$ for $p \neq s$. Thus $u(v^H)$ detects the non-contractibility of v .

4. Let a, b, w, H be as in Example 5.3. The nanoword $v = ww$ lies in the kernel of $\tilde{\gamma}$ and has a zero self-linking function. (Indeed, $R_a = R_b = \mathbb{Z}/2\mathbb{Z}$ and therefore $u(ww) = u(w) + u(w) = 0$ for any nanoword w .) The non-contractibility of v is detected by μ_{v^H} , since $\mu_{v^H}(a, b) = 2\mu_{w^H}(a, b) = 2$.

6.5. Applications. We can use the self-linking function to give a homotopy classification of the monoliteral words a^m where $m \geq 3$ and $a \in \alpha$ with $a \neq \tau(a)$. Desingularizing a^m , we obtain a nanoword $w = (a^m)^d$ of length $m(m-1)$. A direct computation from the definitions gives $[b]_w = 0$ for all $b \in \alpha - \{a\}$ and

$$[a]_w = \sum_{1 \leq i < j \leq m, i+j \neq m+1} a^{(m+1-i-j)(j-i)}.$$

Then $u^w(a) = [a]_w$, $u^w(\tau(a)) = -[a]_w$ and $u^w(b) = 0$ for $b \in \alpha - \{a, \tau(a)\}$. Since $u^w \neq 0$, the word a^m is non-contractible. We can estimate its norm from below applying the last remark of Sect. 6.2 to the monomial determined by $i = 1, j = [m/2] + 1$. This gives

$$(6.5.1) \quad \|a^m\| \geq \left\lfloor \frac{m}{2} \right\rfloor \left\lfloor \frac{m-1}{2} \right\rfloor + 1.$$

(Presumably, $\|a^m\| = m(m-1)/2$.)

Theorem 6.5.1. *Let a, b be letters in α with $a \neq \tau(a), b \neq \tau(b)$. The words a^m and b^n with $m, n \geq 3$ are homotopic if and only if $a = b$ and $m = n$.*

Proof. For $w = a^m$, we can recover both a and m from $u(w) = u(w^d)$. Indeed, a is the only letter on which the value of u^w is a sum of monomials with positive coefficients. The sum of these coefficients is equal to $m(m-1)/2 - [m/2]$. This implies the theorem. $\square$

Let $a \in \alpha$ be a letter with $a \neq \tau(a)$. The self-linking function allows us to show that the word a^m with $m \geq 3$ is not homotopic to a word $v = bcb$ with distinct $b, c \in \alpha$. If $b = \tau(c)$, then v is contractible while a^m is not. If $b \neq \tau(c)$, then the equalities $[b]_v = c, [c]_v = b^{-1}$ imply that the only non-zero values of the self-linking function u^v are $\pm c, \pm b^{-1}$. The sum of coefficients of each of these values is $\pm 1 \neq m(m-1)/2 - [m/2]$. Hence a^m is not homotopic to v .

7. α -FORMS AND α -PAIRINGS

The interlacement of letters in a nanoword $(\mathcal{A}, w)$ is reflected in the function $n_w : \mathcal{A} \times \mathcal{A} \rightarrow \{-1, 0, 1\}$ defined in Sect. 5.1. A study of n_w and related pairings leads to notions of α -forms and α -pairings. We introduce here an abstract algebraic theory of α -forms and α -pairings. Connections with words will be discussed in the next section.

7.1. α -forms. Let $\pi = \pi(\alpha, \tau)$ be the multiplicative abelian group defined in Sect. 5.1. Given a set S , a pairing $l : S \times S \rightarrow \pi$ is *skew-symmetric* if $l(A, B) = l(B, A)^{-1}$ and $l(A, A) = 1$ for all $A, B \in S$. A (skew-symmetric) α -form is an α -alphabet $\mathcal{A}$ endowed with two skew-symmetric pairings $n : \mathcal{A} \times \mathcal{A} \rightarrow \{-1, 0, 1\}$ and $l : \mathcal{A} \times \mathcal{A} \rightarrow \pi$. We view n and l as matrices whose rows and columns are numerated by the letters of $\mathcal{A}$. The skew-symmetry of n means that $n(A, B) = -n(B, A)$ for all $A, B \in \mathcal{A}$ and in particular $n(A, A) = 0$ for all $A \in \mathcal{A}$. Two α -forms $(\mathcal{A}, n, l)$ and $(\mathcal{A}', n', l')$ are *isomorphic* if there is an isomorphism of α -alphabets $\mathcal{A} \rightarrow \mathcal{A}'$ transforming n, l into n', l' , respectively.

Given two α -forms $(\mathcal{A}_1, n_1, l_1)$ and $(\mathcal{A}_2, n_2, l_2)$, we define their direct sum by

$$(\mathcal{A}_1, n_1, l_1) \oplus (\mathcal{A}_2, n_2, l_2) = (\mathcal{A}_1 \amalg \mathcal{A}_2, n_1 \oplus n_2, l_1 \oplus l_2)$$

where $\amalg$ is the disjoint union of α -alphabets and $\oplus$ is the direct sum of matrices. This means that the restriction of $n = n_1 \oplus n_2$ to $\mathcal{A}_i$ is n_i for $i = 1, 2$ and $n(\mathcal{A}_1, \mathcal{A}_2) = n(\mathcal{A}_2, \mathcal{A}_1) = 0$. The restriction of $l = l_1 \oplus l_2$ to $\mathcal{A}_i$ is l_i for $i = 1, 2$ and $l(\mathcal{A}_1, \mathcal{A}_2) = l(\mathcal{A}_2, \mathcal{A}_1) = 1$. Here we assume $\mathcal{A}_1$ and $\mathcal{A}_2$ to be disjoint; if it is not the case we replace $(\mathcal{A}_1, n_1, l_1)$ by an isomorphic α -form and proceed as above. The *trivial α -form* $(\mathcal{A} = \emptyset, n, l)$ yields a neutral element for the direct sum.

For each α -form $f = (\mathcal{A}, n, l)$, we define the *opposite α -form* $f^- = (\mathcal{A}, n^-, l^-)$ where $n^- = -n$ and $l^-(A, B) = (l(A, B))^{-1} = l(B, A)$ for $A, B \in \mathcal{A}$. We also define the *inverse α -form* $\bar{f} = (\bar{\mathcal{A}}, n, l^-)$ where $\bar{\mathcal{A}}$ is the inverse α -alphabet.

7.2. Relation of homology. We define three transformations (moves) on α -forms.

(i)* The move applies to an α -form $(\mathcal{A}, n, l)$ if there is a letter $A \in \mathcal{A}$ such that $n(A, \mathcal{A}) = 0$ and $l(A, \mathcal{A}) = 1$. The move replaces $(\mathcal{A}, n, l)$ with $(\mathcal{A}' = \mathcal{A} - \{A\}, n|_{\mathcal{A}'}, l|_{\mathcal{A}'})$.

(ii)* The move applies to an α -form $(\mathcal{A}, n, l)$ if there are letters $A, B \in \mathcal{A}$ such that $|A| = \tau(|B|)$ and $n(A, C) = n(B, C), l(A, C) = l(B, C)|B|^{n(B, C)}$ for all $C \in \mathcal{A}$. The move replaces $(\mathcal{A}, n, l)$ with $(\mathcal{A}' = \mathcal{A} - \{A, B\}, n|_{\mathcal{A}'}, l|_{\mathcal{A}'})$.

Note that for $C = A$ and $C = B$, the conditions $n(A, C) = n(B, C), l(A, C) = l(B, C)|B|^{n(B, C)}$ are equivalent to $n(A, B) = 0, l(A, B) = 1$. Note also that the formula $l(A, C) = l(B, C)|B|^{n(B, C)}$ is symmetric in A, B . Indeed, this formula is equivalent to $l(B, C) = l(A, C)|A|^{n(A, C)}$ since $|A| = \tau(|B|) = |B|^{-1} \in \pi$ and $n(A, C) = n(B, C)$.

(iii)* The move applies to an α -form $(\mathcal{A}, n, l)$ if there are letters $A, B, C \in \mathcal{A}$ such that $|A| = |B| = |C|$ and $n(A, B) = n(B, C) = 1, n(A, C) = 0$. The move

replaces $(\mathcal{A}, n, l)$ with $(\mathcal{A}, n', l')$ where n', l' coincide with n, l , respectively, except on the pairs $(A, B), (B, A), (A, C), (C, A), (B, C), (C, B)$. The values of n', l' on these pairs are determined by $n'(A, B) = n'(B, C) = 0, n'(A, C) = 1$ and

$$l'(A, B) = l(A, B) |C|, \quad l'(A, C) = l(A, C) |B|^{-1}, \quad l'(B, C) = l(B, C) |A|.$$

Two α -forms are *homologous* if they can be related by a finite sequence of moves (i)*–(iii)*, the inverse moves, and isomorphisms. This defines an equivalence relation of *homology* in the class of α -forms. It is easy to check that α -forms opposite (resp. inverse) to homologous α -forms are themselves homologous.

7.3. α -pairings. To study α -forms, we introduce a simpler concept of an α -pairing. The principal advantage of α -pairings is that for them the move (iii)* becomes trivial.

A (skew-symmetric) α -pairing is a set S endowed with a distinguished element $s \in S$, a mapping $S - \{s\} \rightarrow \alpha$, and a skew-symmetric pairing $b : S \times S \rightarrow \pi = \pi(\alpha, \tau)$. The conditions on S can be rephrased by saying that S is a disjoint union of an α -alphabet and a distinguished element s . As usual, the image of $A \in S - \{s\}$ under the projection to α is denoted $|A|$. Two α -pairings (S, s, b) and (S', s', b') are *isomorphic* if there is a bijection $S \rightarrow S'$ transforming s to s' , inducing an isomorphism of α -alphabets $S - \{s\} \rightarrow S' - \{s'\}$, and transforming b into b' .

The *direct sum* of two α -pairings (S_1, s_1, b_1) and (S_2, s_2, b_2) is the α -pairing

$$(S_1, s_1, b_1) \oplus (S_2, s_2, b_2) = ((S_1 - \{s_1\}) \amalg (S_2 - \{s_2\}) \amalg \{s\}, s, b)$$

where b is uniquely determined by the following conditions: $b(S_1 - \{s_1\}, S_2 - \{s_2\}) = 1$, for $i = 1, 2$, the restriction of b to $S_i - \{s_i\}$ is b_i , and $b(A, s) = b_i(A, s_i)$ for all $A \in S_i - \{s_i\}$. Here we assume S_1 and S_2 to be disjoint; if it is not the case we replace (S_1, s_1, b_1) by an isomorphic α -pairing and proceed as above. The *trivial α -pairing* $(S = \{s\}, s, b = 1)$ yields a neutral element for the direct sum.

For each α -pairing $p = (S, s, b)$, we define the *opposite* α -pairing $p^- = (S, s, b^-)$ where $b^-(A, B) = (b(A, B))^{-1} = b(B, A)$ for $A, B \in S$. We define the *inverse* α -pairing $\bar{p} = (\bar{S}, s, b^-)$ where $\bar{S}$ is the set S with the given mapping $S - \{s\} \rightarrow \alpha$ replaced by its composition with $\tau : \alpha \rightarrow \alpha$.

7.4. Relation of homology for α -pairings. Given an α -pairing (S, s, b) , we say that $A \in S - \{s\}$ is an *annihilating element* if $b(A, C) = 1$ for all $C \in S$. Two elements $A, B \in S - \{s\}$ are *twins* if $|A| = \tau(|B|)$ and $b(A, C) = b(B, C)$ for all $C \in S$. Note that if A, B are twins, then $b(A, B) = b(B, B) = 1$. We define two “compression” moves M_1, M_2 on (S, s, b) . The move M_1 deletes an annihilating element $A \in S - \{s\}$ that is replaces (S, s, b) with $(S' = S - \{A\}, s \in S', b|_{S'})$. The move M_2 deletes a pair of twins $A, B \in S - \{s\}$ that is replaces (S, s, b) with $(S' = S - \{A, B\}, s \in S', b|_{S'})$.

Two α -pairings are *homologous* if they can be related by a finite sequence of moves M_1, M_2 , the inverse moves, and isomorphisms. This defines an equivalence relation of *homology* in the class of α -pairings. It is easy to check that α -pairings opposite (resp. inverse) to homologous α -pairings are themselves homologous.

We now classify α -pairings up to homology. Let us call an α -pairing *primitive* if it has no annihilating elements and no pairs of twins. The following lemma shows that homology classes of α -pairings bijectively correspond to isomorphism classes of primitive α -pairings.

Lemma 7.4.1. *Every α -pairing can be transformed by the moves M_1, M_2 into a primitive α -pairing. Two homologous primitive α -pairings are isomorphic.*

Proof. The first claim is obvious: deleting annihilating elements and pairs of twins by M_1, M_2 we can compress any α -pairing into a primitive one. To prove the second claim, we need the following assertion:

(*) for any $i, j \in \{1, 2\}$, a move M_i^{-1} followed by M_j yields the same result as an isomorphism, or a move M_k , or a move M_k followed by M_l^{-1} with $k, l \in \{1, 2\}$.

This assertion will imply the second claim of the lemma. Indeed, suppose that two primitive α -pairings p, p' are related by a finite sequence of moves $M_1^{\pm 1}, M_2^{\pm 1}$ and isomorphisms. An isomorphism followed by $M_i^{\pm 1}$ can be decomposed as $M_i^{\pm 1}$ followed by an isomorphism. Therefore all isomorphisms in our sequence can be accumulated at the end. The claim (*) implies that p, p' can be related by a finite sequence of moves consisting of several moves of type M_i followed by several moves of type M_i^{-1} and isomorphisms. Since p is primitive we cannot apply to it a move of type M_i . Hence there are no such moves in our sequence. Similarly, since p' (and any isomorphic α -pairing) is primitive, it cannot be obtained by an application of M_i^{-1} . Therefore our sequence consists solely of isomorphisms so that p is isomorphic to p' .

Let us prove (*). We have to consider four cases depending on $i, j \in \{1, 2\}$.

For $i = j = 1$, the move M_i^{-1} on an α -pairing (S, s, b) adds one element A and then M_j removes one element $A' \in S \amalg \{A\}$. If $A' = A$, then $M_j \circ M_i^{-1} = \text{id}$. If $A' \neq A$, then $A' \in S$ is an annihilating element. The transformation $M_j \circ M_i^{-1}$ can be achieved by first applying M_j that removes A' and then applying M_i that adds A .

Let $i = 1, j = 2$. The move M_i^{-1} on (S, s, b) adds an annihilating element A and M_j removes a pair of twins $A_1, A_2 \in S \amalg \{A\}$. If $A_1, A_2 \in S$, then $M_j \circ M_i^{-1}$ can be achieved by first removing A_1, A_2 and then adding A . If $A_1 = A$, then A_2 is an annihilating element of S and $M_j \circ M_i^{-1}$ is the move M_1 removing A_2 . The case $A_2 = A$ is similar.

Let $i = 2, j = 1$. The move M_i^{-1} on (S, s, b) adds a pair of twins A_1, A_2 and M_j removes an annihilating element $A \in S \amalg \{A_1, A_2\}$. If $A \in S$, then $M_j \circ M_i^{-1}$ can be achieved by first removing A and then adding A_1, A_2 . If $A = A_1$, then A_2 is an annihilating element of $S \amalg \{A_2\}$ and $M_j \circ M_i^{-1} = M_1^{-1}$. The case $A = A_2$ is similar.

Let $i = j = 2$. The move M_i^{-1} on (S, s, b) adds a pair of twins A_1, A_2 and M_j removes a pair of twins $A'_1, A'_2 \in S \amalg \{A_1, A_2\}$. If these two pairs are disjoint, then $M_j \circ M_i^{-1}$ can be achieved by first removing $A'_1, A'_2 \in S$ and then adding A_1, A_2 . If these two pairs coincide, then $M_j \circ M_i^{-1} = \text{id}$. It remains to consider the case where these pairs have one common element, say $A'_1 = A_1$, while $A'_2 \neq A_2$. Then $A'_2 \in S$ and for all $C \in S$, $b(A'_2, C) = b(A'_1, C) = b(A_1, C) = b(A_2, C)$. Therefore

the move $M_j \circ M_i^{-1}$ gives an α -pairing isomorphic to (S, s, b) . The isomorphism $S \rightarrow (S - \{A'_2\}) \cup \{A_2\}$ is the identity on $S - \{A'_2\}$ and sends A'_2 into A_2 . $\square$

For an α -pairing p we denote the unique (up to isomorphism) primitive α -pairing homologous to p by p_* . It follows from the definitions that $(p^-)_* = (p_*)^-$, $\overline{p}_* = \overline{p}_*$, and $(p_1 \oplus p_2)_* = (p_1)_* \oplus (p_2)_*$.

7.5. Homology invariants of α -pairings. Any isomorphism invariant of primitive α -pairings yields a homology invariant of an α -pairing $p = (S, s, b)$ by computing on $p_* = (S_*, s_*, b_*)$. For example, $\rho(p) = \text{card}(S_*) - 1$ is a homology invariant of p . Given $a \in \alpha, x \in \pi$, the number

$$\rho_{a,x}(p) = \text{card}\{A \in S_* - \{s_*\} \mid |A| = a, b_*(A, s_*) = x\}$$

is a homology invariant of p . Clearly $\rho(p) = \sum_{a \in \alpha, x \in \pi} \rho_{a,x}(p)$. We have $\rho_{a,x}(p^-) = \rho_{a,x^{-1}}(p)$ and $\rho_{a,x}(\overline{p}) = \rho_{\tau(a),x}(p)$. Both ρ and $\rho_{a,x}$ are additive with respect to the direct sum of α -pairings.

Imitating the self-linking function of nanowords, we can define a homology invariant of $p = (S, s, b)$ as follows. For $a \in \alpha$, set

$$[a]_p = \sum_{A \in S - \{s\}, b(A,s) \neq 1, |A|=a} b(A, s) \in \mathbb{Z}\pi.$$

Define a function $u^p = u(p)$ on α by $u^p(a) = [a]_p \pmod{2} \in (\mathbb{Z}/2\mathbb{Z})\pi$ if $\tau(a) = a$ and $u^p(a) = [a]_p - [\tau(a)]_p \in \mathbb{Z}\pi$ if $\tau(a) \neq a$. It is obvious that u^p is invariant under the moves M_1, M_2 and is thus a homology invariant of p . For $\tau(a) = a$,

$$u^p(a) = u^{p^*}(a) = \sum_{x \in \pi - \{1\}} \rho_{a,x}(p) x \pmod{2}$$

and for $\tau(a) \neq a$,

$$u^p(a) = u^{p^*}(a) = \sum_{x \in \pi - \{1\}} (\rho_{a,x}(p) - \rho_{\tau(a),x}(p)) x.$$

7.6. From α -forms to α -pairings. Every α -form $(\mathcal{A}, n, l)$ determines an α -pairing as follows. Set $S = \mathcal{A} \cup \{s\}$ where s is not a letter of the alphabet $\mathcal{A}$. There is a unique skew-symmetric pairing $b : S \times S \rightarrow \pi$ such that for any $A, B \in \mathcal{A} \subset S$,

$$b(A, s) = \prod_{C \in \mathcal{A}} |C|^{n(A,C)} \in \pi \quad \text{and} \quad b(A, B) = (l(A, B))^2 |A|^{n(A,B)} |B|^{n(A,B)} \in \pi.$$

The α -pairing (S, s, b) associated with $f = (\mathcal{A}, n, l)$ is denoted $p(f)$. The operation $f \mapsto p(f)$ commutes with direct sum and the operations of taking the opposite and the inverse.

Lemma 7.6.1. *α -pairings determined by homologous α -forms are homologous.*

Proof. It suffices to analyze the behavior of the associated α -pairing under the moves (i)*–(iii)* on α -forms. Consider the move (i)* replacing an α -form $(\mathcal{A}, n, l)$ with $(\mathcal{A}' = \mathcal{A} - \{A\}, n|_{\mathcal{A}'}, l|_{\mathcal{A}'})$ where $A \in \mathcal{A}$ is such that $n(A, \mathcal{A}) = 0$ and $l(A, \mathcal{A}) = 1$. Let $(S = \mathcal{A} \cup \{s\}, s, b)$ be the α -pairing associated with $(\mathcal{A}, n, l)$. It

follows from the definitions that A is an annihilating element for b . The α -pairing associated with $(\mathcal{A}', n|_{\mathcal{A}'}, l|_{\mathcal{A}'})$ is $(S' = S - \{A\}, s \in S', b|_{S'})$. It is obtained from (S, s, b) by the move M_1 deleting A .

Consider the move (ii)* replacing $(\mathcal{A}, n, l)$ with $(\mathcal{A}' = \mathcal{A} - \{A, B\}, n|_{\mathcal{A}'}, l|_{\mathcal{A}'})$ where $A, B \in \mathcal{A}$ are such that $|A| = \tau(|B|)$ and $n(A, C) = n(B, C), l(A, C) = l(B, C)|B|^{n(B, C)}$ for all $C \in \mathcal{A}$. Let $(S = \mathcal{A} \cup \{s\}, s, b)$ be the α -pairing associated with $(\mathcal{A}, n, l)$. For $C \in \mathcal{A}$,

$$\begin{aligned} b(A, C) &= (l(A, C))^2 |A|^{n(A, C)} |C|^{n(A, C)} = (l(B, C))^2 |B|^{2n(B, C)} |A|^{n(B, C)} |C|^{n(B, C)} \\ &= (l(B, C))^2 |B|^{n(B, C)} |C|^{n(B, C)} = b(B, C). \end{aligned}$$

Similarly, $b(A, s) = b(B, s)$. Therefore A and B are twins. The α -pairing associated with $(\mathcal{A}', n|_{\mathcal{A}'}, l|_{\mathcal{A}'})$ is $(S' = S - \{A, B\}, s \in S', b|_{S'})$. It is obtained from (S, s, b) by the move M_2 deleting A, B .

Finally, under a move $(\mathcal{A}, n, l) \mapsto (\mathcal{A}, n', l')$ of type (iii)*, the associated α -pairing (S, s, b) is preserved. Indeed, let (S, s, b') be the α -pairing associated with $(\mathcal{A}, n', l')$. It is clear that b' coincides with b except possibly on the pairs $(A, B), (B, A), (A, C), (C, A), (B, C), (C, B)$. We have

$$\begin{aligned} b'(A, B) &= (l'(A, B))^2 |A|^{n'(A, B)} |B|^{n'(A, B)} = (l'(A, B))^2 = (l(A, B)|C|)^2 \\ &= (l(A, B))^2 |A| |B| = (l(A, B))^2 |A|^{n(A, B)} |B|^{n(A, B)} = b(A, B). \end{aligned}$$

Similarly, $b'(B, C) = b(B, C)$ and $b'(A, C) = b(A, C)$. By skew-symmetry, $b' = b$. $\square$

Passing from an α -form to the associated α -pairing we loose certain information contained in the 2-subgroup of π . It is interesting to find further invariants of α -forms retaining this information.

8. LINKING FORMS OF NANOWORDS

We associate with each nanoword w over α an α -form and an α -pairing reflecting the linking of letters in w .

8.1. Notation. Given a nanoword $(\mathcal{A}, w : \hat{n} \rightarrow \mathcal{A})$ and a letter $A \in \mathcal{A}$, we denote by i_A (resp. j_A) the minimal (resp. the maximal) element of the 2-element set $\omega^{-1}(A) \subset \hat{n}$. Thus $w = \cdots A \cdots A \cdots$ where the first entry of A appears on the i_A -th position and the second entry of A appears on the j_A -th position.

8.2. Linking form. For a nanoword $(\mathcal{A}, w : \hat{n} \rightarrow \mathcal{A})$, we define an α -form $f^w = (\mathcal{A}, n_w, lk_w)$ called the *linking form* of w . The function n_w was defined in Sect. 5.1. For letters $D, E \in \mathcal{A}$, set

$$D \circ E = \prod_{F \in \mathcal{A}, i_D < i_F < j_D \text{ and } i_E < j_F < j_E} |F| \in \pi$$

and

$$lk_w(D, E) = (D \circ E)(E \circ D)^{-1} \in \pi.$$

In particular, if $w = \cdots D \cdots D \cdots E \cdots E \cdots$, then $E \circ D = 1$ and $lk_w(D, E) = D \circ E$. If $w = \cdots D \cdots E \cdots E \cdots D \cdots$, then

$$lk_w(D, E) = \prod_{F \in \mathcal{A}, i_D < i_F < i_E < j_F < j_E} |F| \prod_{F \in \mathcal{A}, i_E < i_F < j_E < j_F < j_D} |F|^{-1}.$$

If $w = \cdots D \cdots E \cdots D \cdots E \cdots$, then

$$lk_w(D, E) = \prod_{F \in \mathcal{A}, i_D < i_F < i_E < j_F < j_D \text{ or } i_E < i_F < j_D < j_F < j_E \text{ or } i_D < i_F < i_E < j_D < j_F < j_E} |F|.$$

For all other positions of D, E in w , we can use these formulas to compute $lk_w(E, D)$ and then use the skew-symmetry of lk_w to compute $lk_w(D, E)$.

Lemma 8.2.1. *Under the homotopy moves (i), (ii), (iii) on nanowords, the associated α -forms are transformed by the moves (i)*, (ii)*, (iii)*, respectively.*

Proof. Consider the first homotopy move $w = (\mathcal{A}, xAAy) \mapsto (\mathcal{A} - \{A\}, xy) = v$. Pick $D, E \in \mathcal{A} - \{A\}$. The formulas for $lk_w(D, E)$ show that if a letter $F \in \mathcal{A}$ contributes non-trivially to $lk_w(D, E)$, then $i_F < j_F - 1$. Hence the letter A does not contribute to $lk_w(D, E)$. This implies that lk_v is the restriction of lk_w to $\mathcal{A}'$. It is also obvious that n_v is the restriction of n_w to $\mathcal{A}'$. It remains to observe that $n_w(A, \mathcal{A}) = 0$ and $lk_w(A, \mathcal{A}) = 1$. Thus f^v is obtained from f^w by (i)*.

Consider the move $w = (\mathcal{A}, xAB yBAz) \mapsto (\mathcal{A} - \{A, B\}, xyz) = v$ where $|A| = \tau(|B|)$. It follows from the definitions that $n(A, B) = 0$ and $lk_w(A, B) = 1$. A case by case study of all possible positions of two entries of a letter $C \in \mathcal{A} - \{A, B\}$ in w with respect to AB, BA gives $n(A, C) = n(B, C)$ and $lk_w(A, C) = lk_w(B, C)|B|^{n(B, C)}$. It is obvious that n_v is the restriction of n_w to $\mathcal{A}' = \mathcal{A} - \{A, B\}$. Since the contributions of the letters A, B to $lk_w(C, C')$ cancel each other for all $C, C' \in \mathcal{A} - \{A, B\}$, we obtain that lk_v is the restriction of lk_w to $\mathcal{A}'$. Thus f^v is obtained from f^w by (ii)*.

Consider the move $w = (\mathcal{A}, xAB yAC zBCt) \mapsto (\mathcal{A}, xBA yCA zCBt) = v$ where $|A| = |B| = |C|$. It follows from the definitions that $n_w(A, B) = n_w(B, C) = 1, n_w(A, C) = 0$ and $n_v(A, B) = n_v(B, C) = 0, n_v(A, C) = 1$. It is obvious that for any $D, E \in \mathcal{A} - \{A, B, C\}$, we have $lk_w(D, E) = lk_v(D, E)$. A case by case study of possible positions of the two entries of D in w give

$$lk_w(D, A) = lk_v(D, A), \quad lk_w(D, B) = lk_v(D, B), \quad lk_w(D, C) = lk_v(D, C)$$

for any $D \in \mathcal{A} - \{A, B, C\}$. It remains to observe that

$$lk_v(A, B) = lk_w(A, B)|C|, \quad lk_v(A, C) = lk_w(A, C)|B|^{-1}$$

and $lk_v(B, C) = lk_w(B, C)|A|$. Thus f^v is obtained from f^w by (iii)*. $\square$

This lemma implies that the homology class of the α -form $f^w = (\mathcal{A}, n_w, lk_w)$ is a homotopy invariant of w . This form behaves in a natural way under the standard operations on nanowords:

$$f^{\overline{w}} = \overline{f^w}, \quad f^{w^-} = (f^w)^-, \quad f^{w_1 w_2} = f^{w_1} \oplus f^{w_2}.$$

8.3. Linking pairing. For a nanoword w , the α -pairing associated with the α -form f^w is denoted p^w and called the *linking pairing* of w . It behaves in a natural way under the standard operations on nanowords:

$$p^{\overline{w}} = \overline{p^w}, \quad p^{w^-} = (p^w)^-, \quad p^{w_1 w_2} = p^{w_1} \oplus p^{w_2}.$$

As we know, the α -pairing p^w can be compressed to a primitive α -pairing $p_*^w = (S_*, s_*, b_* : S_* \times S_* \rightarrow \pi)$. The isomorphism class of p_*^w is a homotopy invariant of w . If w is contractible, then p_*^w is a trivial α -form. The formulas above imply that

$$p_*^{\overline{w}} = \overline{p_*^w}, \quad p_*^{w^-} = (p_*^w)^-, \quad p_*^{w_1 w_2} = p_*^{w_1} \oplus p_*^{w_2}.$$

The homology invariants of α -pairings defined in Sect. 7.5 yield homotopy invariants of nanowords. In particular, set $\rho(w) = \rho(p^w) = \text{card}(S_*) - 1$ and for $a \in \alpha, x \in \pi$, set

$$\rho_{a,x}(w) = \rho_{a,x}(p^w) = \text{card}\{A \in S_* - \{s_*\} \mid |A| = a, b_*(A, s_*) = x\}.$$

We have $\rho_{a,x}(\overline{w}) = \rho_{\tau(a),x}(w)$, $\rho_{a,x}(w^-) = \rho_{a,x^{-1}}(w)$ and $\rho_{a,x}(w_1 w_2) = \rho_{a,x}(w_1) + \rho_{a,x}(w_2)$. Application: if $\rho_{a,x}(w) \neq 0$ for some a, x , then ww' is non-contractible for any nanoword w' .

For any nanoword $(\mathcal{A}, w)$, the homotopy invariance of ρ implies that $\|w\| \geq \rho(w)$. Application: if the α -pairing p^w is primitive, then w is not homotopic to a nanoword of smaller length and $\|w\| = \rho(w) = \text{card}(\mathcal{A})$.

Note finally that the self-linking function u^w of a nanoword w is computed from p_*^w : it follows from the definitions that $u^w = u(p^w) = u(p_*^w)$.

8.4. Nanowords of length 4. It follows from the definitions that all nanowords of length ≤ 2 are contractible. We give a homotopy classification of nanowords of length 4. Clearly, each such nanoword is either contractible or isomorphic to the nanoword $(\mathcal{A} = \{A, B\}, w = ABAB)$ with $|A| = a \in \alpha, |B| = b \in \alpha$. Denote the latter nanoword by $w_{a,b}$ (possibly $a = b$). Clearly, $\overline{w_{a,b}} = w_{\tau(a),\tau(b)}$ and $(w_{a,b})^- \approx w_{b,a}$.

By the second homotopy move, if $b = \tau(a)$, then $w_{a,b}$ is contractible. The following theorem establishes the converse and gives a homotopy classification of these nanowords.

Theorem 8.4.1. *For $a, b \in \alpha$ with $\tau(a) \neq b$, the nanoword $w_{a,b}$ is non-contractible. Two nanowords $w_{a,b}$ and $w_{a',b'}$ with $\tau(a) \neq b, \tau(a') \neq b'$ are homotopic if and only if $a = a'$ and $b = b'$.*

Proof. Set $w = w_{a,b}$. The linking α -form $(\mathcal{A}, n_w, lk_w)$ is given by the matrices

$$n_w = \begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix}, \quad lk_w = \begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix}.$$

Here the 1-st and 2-nd rows (and columns) correspond to A, B , respectively. The associated α -pairing $p^w = (S, s, S \times S \rightarrow \pi)$ with $S = \{s, A, B\}$ is given by the

matrix

$$(8.4.1) \quad p^w = \begin{bmatrix} 1 & b^{-1} & a \\ b & 1 & ab \\ a^{-1} & a^{-1}b^{-1} & 1 \end{bmatrix}.$$

Here the rows (resp. columns) correspond to $s, A, B \in S$ counting from top to bottom (resp. from left to right). Since $b \neq \tau(a)$, we have $ab \neq 1 \in \pi$ so that p^w has no annihilating elements and no twins. Hence p^w is primitive and $\|w\| = \rho(w) = 2$.

Suppose that the nanowords $w = ABAB$ and $w' = A'B'A'B'$ with $|A| = a, |B| = b \neq \tau(a), |A'| = a', |B'| = b' \neq \tau(a')$ are homotopic. Then the associated primitive α -pairings are isomorphic. If the isomorphism sends A, B to A', B' , respectively, then $a' = a, b' = b$ and we are done. If the isomorphism sends A to B' and B to A' , then $a' = b$ and $b' = a$. We claim that in the latter case $a = b$. If $a \neq b$, then a, b lie in different orbits $\hat{a}, \hat{b} \subset \alpha$ of τ . Recall from Sect. 4.2 the functions $\mu_w, \mu_{w'} : \hat{\alpha} \times \hat{\alpha} \rightarrow \mathbb{Z}$ where $\hat{\alpha}$ is the set of orbits of τ . Clearly, $\mu_w(\hat{a}, \hat{b}) = 1$ and $\mu_{w'}(\hat{a}, \hat{b}) = -\mu_{w'}(\hat{b}, \hat{a}) = -1$. This contradicts the invariance of μ_w under homotopy. Therefore $a = b = a' = b'$. $\square$

Corollary 8.4.2. *A nanoword of length ≤ 4 is homotopically skew-symmetric if and only if it is contractible. A nanoword of length ≤ 4 is homotopically symmetric if and only if it is contractible or symmetric.*

Proof. The nanoword $w = w_{a,b}$ is homotopically skew-symmetric if and only if $w \simeq w_{\tau(b), \tau(a)}$. By Theorem 8.4.1, this happens if and only if $\tau(a) = b$. The nanoword $w = w_{a,b}$ is homotopically symmetric if and only if $w \simeq w_{b,a}$. By Theorem 8.4.1, this happens if and only if $\tau(a) = b$ or $a = b$. $\square$

Corollary 8.4.3. *The group of homotopy automorphisms of $w_{a,b}$ is the full group of τ -equivariant bijections $\alpha \rightarrow \alpha$ if $a = \tau(b)$ and its subgroup consisting of bijections preserving a and b if $a \neq \tau(b)$.*

8.5. Example. The following example shows that for the nanowords of length 6, the associated α -pairing is a strong but non-faithful invariant. Pick three letters $a, b, c \in \alpha$ (possibly coinciding) and consider the α -alphabet $\mathcal{A} = \{A, B, C\}$ where $|A| = a, |B| = b, |C| = c$. Consider the nanoword $(\mathcal{A}, w = ABACBC)$. The linking α -form $(\mathcal{A}, n_w, lk_w)$ is given by

$$n_w = \begin{bmatrix} 0 & 1 & 0 \\ -1 & 0 & 1 \\ 0 & -1 & 0 \end{bmatrix}, \quad lk_w = \begin{bmatrix} 1 & 1 & b \\ 1 & 1 & 1 \\ b^{-1} & 1 & 1 \end{bmatrix}.$$

Here the rows and columns correspond to A, B, C , respectively. For example, $lk_w(A, B) = 1 \in \pi, lk_w(A, C) = b \in \pi$, etc. The associated α -pairing $p^w = (S, s, S \times S \rightarrow \pi)$ with $S = \{s, A, B, C\}$ is given by

$$p^w = \begin{bmatrix} 1 & b^{-1} & ac^{-1} & b \\ b & 1 & ab & b^2 \\ a^{-1}c & a^{-1}b^{-1} & 1 & bc \\ b^{-1} & b^{-2} & b^{-1}c^{-1} & 1 \end{bmatrix}.$$

Here the rows and columns correspond to s, A, B, C , respectively.

Observe that all the generators $\{e\}_{e \in \alpha}$ of π represent non-trivial elements of π . More generally, a product of an odd number of generators is a non-trivial element of π . This implies that $b^{\pm 1} \neq a^{-1}c$ so that A, B are not twins and B, C are not twins. It is also clear that A and C are not annihilating elements. The element $B \in S$ is annihilating if and only if $a = c = \tau(b)$. The letters A and C are twins if and only if $a = \tau(c)$ and $b = \tau(b)$. In all other cases p^w is primitive and $\|w\| = 3$. If $a = c = \tau(b)$, then deleting B we obtain a homologous α -pairing

$$\begin{bmatrix} 1 & a & a^{-1} \\ a^{-1} & 1 & a^{-2} \\ a & a^2 & 1 \end{bmatrix}.$$

If $a \neq \tau(a)$, then this α -pairing is primitive and $3 \geq \|w\| \geq \rho(w) = 2$. If $a = b = c = \tau(a)$, then w is contractible by $w \simeq BACACB \simeq BB \simeq \emptyset$. If $a = \tau(c)$ and $b = \tau(b)$, then deleting A, C we obtain a homologous α -pairing

$$\begin{bmatrix} 1 & a^2 \\ a^{-2} & 1 \end{bmatrix}.$$

If $a \neq \tau(a)$, then it is primitive and $3 \geq \|w\| \geq \rho(w) = 1$. In the remaining case $a = \tau(a) = c \neq b = \tau(b)$, the α -pairing p^w is homologous to a trivial α -pairing and gives no information about w . The homomorphisms γ and $\tilde{\gamma}$ do not help since in this case $\tilde{\gamma}(w) = 1$. However, w is non-contractible by Theorem 13.2.1 below.

The nanoword $ABACBC$ with $|A| = |B| = |C| = c$ is the desingularization of the word ccc in the alphabet α . Therefore if $c \neq \tau(c)$, then

$$(8.5.1) \quad \|ccc\| = 3.$$

9. ANALYSIS LITTERAE

We give a homotopy classification of words of length ≤ 5 . Since the desingularization of a word deletes all letters of multiplicity 1, we consider only words which contain no letters of multiplicity 1. We call such words *multiplicity-one-free*. We first summarize the results concerning the words of length ≤ 4 .

Theorem 9.0.1. *A multiplicity-one-free word of length ≤ 4 in the alphabet α has one of the following forms: $aa, a^3 = aaa, a^4 = aaaa, aabb, abba, bbaa, abab$ with distinct $a, b \in \alpha$. The words $aa, aabb, abba, bbaa$ are contractible. The words a^3, a^4 are contractible if and only if $\tau(a) = a$. The word $abab$ is contractible if and only if $\tau(a) = b$. Non-contractible words of type $a^3, a^4, abab$ are homotopic if and only if they are equal, i.e., coincide letter-wise.*

Proof. That all multiplicity-one-free words of length ≤ 4 belong to our list is obvious. The words $aa, aabb, abba, bbaa$ are nanowords and can be contracted by the first homotopy move. If $\tau(a) = a$, then a^3, a^4 are contractible by Example 3.4.3. If $\tau(a) \neq a$, then a^3, a^4 are not contractible by Sect. 6.5. If $\tau(a) = b \neq a$, then $abab$ is contractible by the second homotopy move. If $\tau(a) \neq b \neq a$, then $abab$ is not contractible by Theorem 8.4.1. The same theorem shows that two

non-contractible homotopic words of this type are equal. That non-contractible homotopic words of type a^3 or a^4 are equal follows from Theorem 6.5.1. That a word $abab$ with $\tau(a) \neq b \neq a$ is not homotopic to a monoliteral word follows from the results of Sect. 6.5. $\square$

Theorem 9.0.2. *A multiplicity-one-free word of length 5 in the alphabet α has one of the following forms:*

$$(9.0.2) \quad aaabb, aabba, abbaa, bbaaa,$$

$$(9.0.3) \quad a^5, abaab, baaba, aabab, babaa, baaab, ababa$$

with distinct $a, b \in \alpha$. The words (9.0.2) are homotopic to aaa ; they are contractible if and only if $\tau(a) = a$. The words a^5 and $baaab$ are contractible if and only if $\tau(a) = a$. The word $ababa$ is contractible if and only if $\tau(a) = b$. The words $abaab, baaba, aabab, babaa$ are not contractible. Two non-contractible words of type (9.0.3) are homotopic if and only if they coincide letter-wise with the following exception:

$$aabab \simeq babaa \simeq baaab$$

for $a = \tau(b) \neq b$. Non-contractible words of type (9.0.3) are not homotopic to words of length ≤ 4 .

Note a subtlety in the last claim: it does not exclude that a word of type (9.0.3) is homotopic to a *nanoword* over α of length ≤ 4 (cf. Example 3.4.5). This nanoword however can not be a word in the alphabet α .

Among the words listed in Theorem 9.0.2 only the words $a^5, ababa, baaab$ are symmetric. Here is a complete list of homotopically symmetric words of length 5.

Corollary 9.0.3. *A multiplicity-one-free word of length 5 is homotopically symmetric if and only if it is contractible or symmetric or belongs to the list (9.0.2) or has the form $aabab$ or $babaa$ with $b = \tau(a) \neq a$.*

Among the words listed in Theorem 9.0.2 only the words $a^5, ababa, baaab$ may be skew-symmetric and this happens when $a = \tau(a)$ and $b = \tau(b)$.

Corollary 9.0.4. *A multiplicity-one-free word of length 5 is homotopically skew-symmetric if and only if it is contractible or skew-symmetric.*

Corollary 9.0.5. *Let $\tau = \text{id} : \alpha \rightarrow \alpha$. A multiplicity-one-free word of length ≤ 5 in the alphabet α is non-contractible if and only if it has one of the following six forms:*

$$abab, abaab, baaba, aabab, babaa, ababa$$

where $a \neq b$. Such words are homotopic if and only if they coincide letter-wise.

The rest of this section is devoted to a proof of Theorem 9.0.2.

9.1. Reduction to a lemma. That all multiplicity-one-free words of length 5 belong to our list is obvious. The words (9.0.2) are homotopic to aaa by the first homotopy move. By Example 3.4.3 and Theorem 6.5.1, the words a^3 and a^5 are contractible if and only if $\tau(a) = a$. By Sect. 6.5, if a^5 is non-contractible, then it is not homotopic to a word of length ≤ 4 . In the rest of the proof we use the notation $w_{a,b}^1 = abaab$ and similarly

$$w_{a,b}^2 = baaba, w_{a,b}^3 = aabab, w_{a,b}^4 = babaa, w_{a,b}^5 = baaab, w_{a,b}^6 = ababa.$$

where $a \neq b$. Using Example 3.4.3, it is easy to show that the word $w_{a,b}^5$ with $\tau(a) = a$ is contractible. In the sequel, writing $w_{a,b}^5$ we always suppose that $\tau(a) \neq a$. By Example 3.4.4, the word $w_{a,b}^6$ with $a = \tau(b)$ is contractible. In the sequel, writing $w_{a,b}^6$ we always suppose that $a \neq \tau(b)$.

It remains to prove the following lemma for $i = 1, \dots, 6$.

Lemma 9.1.1. *(*) The word $w_{a,b}^i$ is not homotopic to a word of length ≤ 4 or to a monoliteral word. In particular, $w_{a,b}^i$ is non-contractible.*

*(**) The words $w_{a,b}^i, w_{a',b'}^i$ are homotopic if and only if $a = a'$ and $b = b'$.*

*(***) For $j < i$, the words of type w^i are not homotopic to the words of type w^j with the following exception: $w_{a,b}^3 \simeq w_{a,b}^4 \simeq w_{a,b}^5$ for $a = \tau(b) \neq b$.*

We prove this lemma case by case. The cases $i = 3$ and $i = 6$ need additional techniques at least for some a, b . The proof in this cases will be accomplished in Sect. 12.

9.2. Case $i = 1$. The étale word associated with $w_{a,b}^1 = abaab$ is the pair $(\mathcal{A} = \alpha, aabab)$. Its desingularization yields the nanoword $w = A_3A_2BA_3A_1A_2A_1B$ in the α -alphabet

$$(9.2.1) \quad \{A_1 = a_{2,3}, A_2 = a_{1,3}, A_3 = a_{1,2}, B = b_{1,2}\}$$

where $|A_1| = |A_2| = |A_3| = a, |B| = b$. The linking α -form of w is given by

$$n_w = \begin{bmatrix} 0 & -1 & 0 & 0 \\ 1 & 0 & -1 & 1 \\ 0 & 1 & 0 & 1 \\ 0 & -1 & -1 & 0 \end{bmatrix}, \quad lk_w = \begin{bmatrix} 1 & 1 & a^{-1} & 1 \\ 1 & 1 & 1 & a \\ a & 1 & 1 & a \\ 1 & a^{-1} & a^{-1} & 1 \end{bmatrix}$$

where the rows and columns correspond to A_1, A_2, A_3, B , respectively. Consider the associated α -pairing $p^w = (S, s, S \times S \rightarrow \alpha)$. Here $S = \{s, A_1, A_2, A_3, B\}$ and $S \times S \rightarrow \alpha$ is given by the matrix

$$\begin{bmatrix} 1 & a & b^{-1} & a^{-1}b^{-1} & a^2 \\ a^{-1} & 1 & a^{-2} & a^{-2} & 1 \\ b & a^2 & 1 & a^{-2} & a^3b \\ ab & a^2 & a^2 & 1 & a^3b \\ a^{-2} & 1 & a^{-3}b^{-1} & a^{-3}b^{-1} & 1 \end{bmatrix}$$

where the rows and columns correspond to s, A_1, A_2, A_3, B , respectively. An easy inspection (using only that $a \neq b$) shows that p^w is primitive and therefore $\|w_{a,b}^1\| = \|w\| = 4$. In particular $w_{a,b}^1$ is non-contractible. We can recover a and b

from p^w as follows. Since p^w is primitive, it is determined by the homotopy class of $w_{a,b}^1$ uniquely up to isomorphism. The projection $S - \{s\} \rightarrow \alpha$ takes one value (the letter a) with multiplicity 3 and another value (the letter b) with multiplicity 1. This “multiplicity argument” shows that if $w_{a,b}^1 \simeq w_{a',b'}^1$, then $a = a', b = b'$. The equality $\|w_{a,b}^1\| = 4$ implies that $w_{a,b}^1$ is not homotopic to a nanoword of length ≤ 4 . That w is not homotopic to a monoliteral word follows from the fact that the projection $S - \{s\} \rightarrow \alpha$ takes two distinct values while the similar projection associated with a monoliteral word takes only one value.

9.3. Case $i = 2$. Since $w_{a,b}^2$ is opposite to $w_{a,b}^1$, we have $\|w_{a,b}^2\| = \|w_{a,b}^1\| = 4$ and the claims $(*)$, $(**)$ for $i = 1$ imply similar claims for $i = 2$. We need to check only that $w_{a,b}^2$ is not homotopic to $w_{a',b'}^1$. As we know, the α -pairing of $w_{a,b}^2 = (w_{a,b}^1)^-$ is opposite to the one of $w_{a,b}^1$. If $w_{a,b}^2 \simeq w_{a',b'}^1$, then the multiplicity argument counting the values of the projection $S - \{s\} \rightarrow \alpha$ shows that $a = a'$ and $b = b'$. The matrix of the α -pairing of $w_{a,b}^2$ is obtained by transposition from the one of $w_{a,b}^1$ (and possibly by simultaneous permutation of rows and columns keeping the first row and the first column). Comparing the values in the first column we obtain that $\{a^{-1}, b\} = \{a, b^{-1}\}$. Since $a \neq b$, we have $\tau(a) = a, \tau(b) = b$. Let $H = \{1, b\} \subset \pi$. The H -coverings of $w_{a,b}^1 \simeq A_3A_2BA_3A_1A_2A_1B$ and $w_{a,b}^2 \simeq BA_3A_2A_3A_1BA_2A_1$ are the nanowords A_2BA_2B and BA_2BA_2 , respectively. Since $a \neq b$, these nanowords are homotopic only if $a = \tau(b)$ which is excluded by $\tau(b) = b$.

9.4. Case $i = 3$. The desingularization of the étale word associated with $w_{a,b}^3 = aabab$ yields the nanoword $A_3A_2A_3A_1BA_2A_1B$ in the α -alphabet (9.2.1). Its linking α -form is given by

$$n = \begin{bmatrix} 0 & -1 & 0 & 1 \\ 1 & 0 & -1 & 1 \\ 0 & 1 & 0 & 0 \\ -1 & -1 & 0 & 0 \end{bmatrix}, \quad lk = \begin{bmatrix} 1 & 1 & a^{-1} & 1 \\ 1 & 1 & 1 & a \\ a & 1 & 1 & a \\ 1 & a^{-1} & a^{-1} & 1 \end{bmatrix}.$$

The associated α -pairing is given by the matrix

$$p = \begin{bmatrix} 1 & ab^{-1} & b^{-1} & a^{-1} & a^2 \\ a^{-1}b & 1 & a^{-2} & a^{-2} & ab \\ b & a^2 & 1 & a^{-2} & a^3b \\ a & a^2 & a^2 & 1 & a^2 \\ a^{-2} & a^{-1}b^{-1} & a^{-3}b^{-1} & a^{-2} & 1 \end{bmatrix}$$

where the rows and columns correspond to s, A_1, A_2, A_3, B , respectively.

We first study the case where $a \neq \tau(b)$ (and $a \neq b$). Then $ab \neq 1$ and p is primitive. The arguments as above show that: $\|w_{a,b}^3\| = 4$; $w_{a,b}^3$ is not homotopic to a nanoword of length ≤ 4 or to a monoliteral word; if $w_{a,b}^3 \simeq w_{a',b'}^3$ with $a' \notin \{b', \tau(b')\}$, then $a = a', b = b'$. If $w_{a,b}^3 \simeq w_{a',b'}^1$ with $a' \neq b'$, then the

multiplicity argument gives $a = a', b = b'$ and comparing the values in the s -column we obtain that $\{a^{-1}, b\} = \{a, b\}$. Hence $a = \tau(a) \neq b$. We must prove the following.

Lemma 9.4.1. *If $a = \tau(a) \neq b$, then the nanowords $A_3A_2A_3A_1BA_2A_1B$ and $A_3A_2BA_3A_1A_2A_1B$ in the α -alphabet (9.2.1) are not homotopic.*

The α -pairings of the nanowords in question are isomorphic and coverings do not help. We prove this lemma in Sect. 12.4 using other techniques.

If $w_{a,b}^3 \simeq w_{a',b'}^2$ with $a' \neq b'$, then as above $a = a', b = b'$ and $\{a, b^{-1}\} = \{a, b\}$, $\{a^{-1}b^{-1}, a^2\} = \{a^{-1}b, a^{-2}\}$. Hence $b = \tau(b) \neq a = \tau(a)$. Let $H = \{1, b\} \subset \pi$. The H -coverings of $w_{a,b}^3$ and $w_{a,b}^2$ are the nanowords A_2BA_2B and BA_2BA_2 . If they are homotopic, then $a = b$ which contradicts the assumptions.

We now study the case $a = \tau(b)$. By Example 3.4.5, $w_{a,b}^3$ is homotopic to the nanoword $w_{a,a}$ of length 4. The latter is non-contractible since $a \neq b = \tau(a)$. Therefore $\|w_{a,b}^3\| = 2$. The word $w_{a,b}^3 \simeq w_{a,a}$ is not homotopic to a word $cdcd$ with distinct $c, d \in \alpha$ by Theorem 8.4.1. The word $w_{a,b}^3$ is not homotopic to a monoliteral word c^m with $c \in \alpha, c \neq \tau(c), m \geq 3$ because $\|c^m\| \geq 3$ by (6.5.1) and (8.5.1). The word $w_{a,b}^3$ is not homotopic to $w_{a',b'}^i$ for $i = 1, 2$ since they have different norms. If $w_{a,b}^3 \simeq w_{a',b'}^3$, then $\|w_{a',b'}^3\| = \|w_{a,b}^3\| = 2$ and we must have $a' = \tau(b')$. Then $w_{a,a} \simeq w_{a',a'}$ and therefore $a = a'$ and $b = \tau(a) = \tau(a') = b'$.

9.5. Case $i = 4$. Since $w_{a,b}^4 = (w_{a,b}^3)^-$, the claims $(*)$, $(**)$ for $i = 4$ follow from the similar claims for $i = 3$. The claim $(***)$ for $i = 3$ implies that $w_{a,b}^4$ is not homotopic to $w_{a',b'}^j$ with $j = 1, 2$. If $a = \tau(b)$, then $w_{a,b}^4 \simeq w_{a,b}^3$ by Example 3.4.5. It remains to prove that $w_{a,b}^4$ with $a \neq \tau(b)$ (and $a \neq b$) is not homotopic to $w_{a',b'}^3$. If they are homotopic then the multiplicity argument yields $a = a'$ and $b = b'$. The matrix of the α -pairing of $w_{a,b}^4 = (w_{a,b}^3)^-$ is obtained by transposition from the one of $w_{a,b}^3$. Comparing the values in the s -column we obtain that $\{a^{-1}, b^{-1}\} = \{a, b\}$. Since $a \neq \tau(b) = b^{-1}$, we have $a = a^{-1} = \tau(a)$ and $b = \tau(b)$. Let $H = \{1, b\} \subset \pi$. The nanowords $(w_{a,b}^3)^H = A_2BA_2B$ and $(w_{a,b}^4)^H = BA_2BA_2$ are homotopic only if $a = \tau(b)$ or $a = b$ which is excluded by the assumptions.

9.6. Case $i = 5$. The desingularization of the étale word associated with $w_{a,b}^5 = baaab$ yields the nanoword $BA_3A_2A_3A_1A_2A_1B$ in the α -alphabet (9.2.1). Its linking α -form is given by

$$n = \begin{bmatrix} 0 & -1 & 0 & 0 \\ 1 & 0 & -1 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix}, \quad lk = \begin{bmatrix} 1 & 1 & a^{-1} & a^{-1} \\ 1 & 1 & 1 & 1 \\ a & 1 & 1 & a \\ a & 1 & a^{-1} & 1 \end{bmatrix}.$$

The associated α -pairing is given by the matrix

$$p = \begin{bmatrix} 1 & a & 1 & a^{-1} & 1 \\ a^{-1} & 1 & a^{-2} & a^{-2} & a^{-2} \\ 1 & a^2 & 1 & a^{-2} & 1 \\ a & a^2 & a^2 & 1 & a^2 \\ 1 & a^2 & 1 & a^{-2} & 1 \end{bmatrix}$$

where the rows and columns correspond to s, A_1, A_2, A_3, B , respectively. By our assumptions $\tau(a) \neq a \neq b$. Assume first that $a \neq \tau(b)$. Then p is primitive and the arguments as above show that: $\|w_{a,b}^5\| = 4$; $w_{a,b}^5$ is not homotopic to a nanoword of length ≤ 4 or to a monoliteral word; if $w_{a,b}^5 \simeq w_{a',b'}^5$ with $a' \notin \{b', \tau(b')\}$, then $a = a', b = b'$. If $w_{a,b}^5 \simeq w_{a',b'}^j$ with $j = 1, 2, 3, 4$, then comparing the s -columns of the corresponding α -pairings we always obtain that $b' = a'$, a contradiction.

We now study the case $a = \tau(b) \neq b$. By Example 3.4.5, $w_{a,b}^5 \simeq w_{a,a}$. As in the case $i = 3$, we obtain that $\|w_{a,b}^5\| = 2$ and $w_{a,b}^5$ verifies the claim $(*)$ of the lemma. If $w_{a,b}^5 \simeq w_{a',b'}^j$ for $j \leq 5$, then comparing the norms we observe that $j \in \{3, 4, 5\}$ and $a' = \tau(b')$. Then $w_{a,a} \simeq w_{a,b}^5 \simeq w_{a',b'}^j \simeq w_{a',a'}$ so that $a = a'$ and $b = \tau(a) = \tau(a') = b'$. Thus we recover either one of the exceptional homotopies from $(**)$ or the tautological homotopy $w_{a,b}^5 \simeq w_{a,b}^5$.

9.7. Case $i = 6$. The desingularization of the étale word associated with $w_{a,b}^6 = ababa$ yields the nanoword $w = A_3A_2BA_3A_1BA_2A_1$ in the α -alphabet (9.2.1). The linking α -form of w is given by

$$n_w = \begin{bmatrix} 0 & -1 & 0 & -1 \\ 1 & 0 & -1 & 0 \\ 0 & 1 & 0 & 1 \\ 1 & 0 & -1 & 0 \end{bmatrix}, \quad lk_w = \begin{bmatrix} 1 & b^{-1} & a^{-1}b^{-1} & 1 \\ b & 1 & b^{-1} & 1 \\ ab & b & 1 & 1 \\ 1 & 1 & 1 & 1 \end{bmatrix}.$$

The associated α -pairing is given by the matrix

$$(9.7.1) \quad p^w = \begin{bmatrix} 1 & ab & 1 & a^{-1}b^{-1} & 1 \\ a^{-1}b^{-1} & 1 & a^{-2}b^{-2} & a^{-2}b^{-2} & a^{-1}b^{-1} \\ 1 & a^2b^2 & 1 & a^{-2}b^{-2} & 1 \\ ab & a^2b^2 & a^2b^2 & 1 & ab \\ 1 & ab & 1 & a^{-1}b^{-1} & 1 \end{bmatrix}$$

where the rows and columns correspond to s, A_1, A_2, A_3, B , respectively. This matrix depends only on $ab \in \pi$. The assumption $a \neq \tau(b)$ guarantees that $ab \neq 1$. It is clear that A_1, A_3, B are not annihilating and A_2 is annihilating if and only if $a^2b^2 = 1$. The latter is equivalent to $a = \tau(a)$ and $b = \tau(b)$. The only possible twins are A_1, A_3 and again this happens if and only if $a = \tau(a), b = \tau(b)$. Thus, if $a = \tau(a), b = \tau(b)$, then p^w is homologous to a trivial α -pairing. Since this is not the case for the words considered above in this lemma (i.e., the words $w^j, j = 1, \dots, 5$, the non-contractible monoliteral words, and the non-contractible nanowords of length 4), w is not homotopic to either of them. It remains to prove the following.

Lemma 9.7.1. *Let $a = \tau(a) \neq b = \tau(b)$. Then $w_{a,b}^6$ is non-contractible. If $w_{a,b}^6 \simeq w_{a',b'}^6$ with $a' = \tau(a') \neq b' = \tau(b')$, then $a = a'$ and $b = b'$.*

We shall prove this lemma in Sect. 12.5.

Consider now the case where $a \neq \tau(a)$ or $b \neq \tau(b)$. Then p^w is primitive and $\|w\| = 4$. The arguments given in the case $i = 1$ apply here and yield the claims $(*)$, $(**)$ for $i = 6$. That $w_{a,b}^6$ is not homotopic to $w_{a',b'}^j$ with $j \leq 5$ follows from the fact that their α -pairings are not isomorphic: the s -column of (9.7.1) does not contain generators $\{e\}_{e \in \alpha} \subset \pi$ while the s -column of the α -pairings associated with w^j contains such generators.

10. COLORINGS OF NANOWORDS

Throughout this section we fix a set $\beta \subset \alpha$ such that $\tau(\beta) = \beta$. (The set β may be empty or equal to α .) We shall use notation introduced in Sect. 8.1.

10.1. Tricolorings. A *tricoloring* (with respect to $\beta \subset \alpha$) of a nanoword $(\mathcal{A}, w : \hat{n} \rightarrow \mathcal{A})$ over α is a function $f : \{0, 1, \dots, n\} \rightarrow \mathbb{Z}/3\mathbb{Z}$ such that for any $A \in \mathcal{A}$ with $|A| \in \beta$,

$$f(i_A) = f(i_A - 1), \quad f(j_A - 1) + f(j_A) + f(i_A) = 0$$

and for any $A \in \mathcal{A}$ with $|A| \in \alpha - \beta$,

$$f(i_A - 1) + f(i_A) + f(j_A) = 0, \quad f(j_A) = f(j_A - 1).$$

Note that a sum of three residues (mod 3) is 0 if and only if these residues are either equal to each other or form the triple $\{0, 1, 2\}$. It is convenient to insert dashes between consecutive letters of w and also at the beginning and the end of w . Thus instead of $w(1)w(2) \cdots w(n)$ we write $-w(1) - w(2) - \cdots - w(n) -$. The residue $f(i)$ can be interpreted as the label of the dash between $w(i)$ and $w(i+1)$. The residues $f(0)$ and $f(n)$ are the labels of the leftmost and the rightmost dashes, respectively. They are called the *input* and the *output* of f , respectively.

The sum of two tricolorings of a nanoword w is a tricoloring of w . Thus the tricolorings of w form a vector space over $\mathbb{Z}/3\mathbb{Z}$. It is non-zero since w has constant tricolorings $f = \text{const}$. Let $c_{k,l} = c(\beta, w)_{k,l}$ be the number of tricolorings of w with given input and output $k, l \in \mathbb{Z}/3\mathbb{Z}$. The difference between two tricolorings with the same input and output has input and output zero. Therefore either $c_{k,l} = 0$ or $c_{k,l} = c_{0,0}$. Adding constant tricolorings and negating, we observe that $c_{k,l}$ depends only on $k - l$ and $c_{k,l} = c_{-k,-l}$. We conclude that the matrix $(c_{k,l})$ has the form

$$(10.1.1) \quad \begin{bmatrix} c & c & c \\ c & c & c \\ c & c & c \end{bmatrix} \quad \text{or} \quad \begin{bmatrix} c & 0 & 0 \\ 0 & c & 0 \\ 0 & 0 & c \end{bmatrix}$$

where $c = c_{0,0} \geq 1$ is a power of 3.

An easy check shows that the number $c(\beta, w)_{k,l}$ is preserved under homotopy moves on w for all k, l , cf. Sect. 10.4. This can be used to distinguish homotopy

types of nanowords (see examples below). In particular, if a nanoword admits non-constant tricolorings, then it is non-contractible.

The formula $w \mapsto (c(\beta, w)_{k,l})_{k,l \in \mathbb{Z}/3\mathbb{Z}}$ defines a representation of the monoid of homotopy classes of nanowords $\mathcal{N}_\bullet(\alpha)$ in the monoid of integral (3×3) -matrices of type (10.1.1). For $\beta = \emptyset$ or $\beta = \alpha$, any nanoword has only constant tricolorings.

10.2. Examples. 1. Consider the nanoword $w = ABAB$ such that $|A|, |B|$ lie in different orbits of α . Take as β the orbit of $|A|$. Then w admits a non-constant tricoloring

$$\begin{array}{cccc} 0 & 0 & 2 & 1 \\ - & A & - B & - A & - B & - \end{array}.$$

This gives another proof of the fact that this nanoword is not contractible.

2. Consider the nanoword $w = A_1 A_2 B A_3 A_1 B A_2 A_3$ where $|A_1| = |A_2| = |A_3|$ and $|B|$ lie in different orbits of τ . Take as β the orbit of $|A_1|$. An easy check shows that $c(\beta, w)_{k,l} = 3$ for all k, l . Here is an example of a coloring of w with distinct input and output:

$$\begin{array}{cccccccc} 0 & 0 & 0 & 1 & 1 & 2 & 2 & 1 & 1 \\ - & A_1 & - A_2 & - B & - A_3 & - A_1 & - B & - A_2 & - A_3 & - \end{array}.$$

Hence w is non-contractible. Note that $w \in \text{Ker}(\tilde{\gamma})$.

10.3. Ring Λ . To define more general colorings, we introduce a group Ψ_α generated by $\{a, a_\bullet\}_{a \in \alpha}$ with defining relations $aa_\bullet = a_\bullet a$ and $a\tau(a) = a_\bullet\tau(a)_\bullet = 1$ for $a \in \alpha$. Let $\Lambda = \mathbb{Z}\Psi_\alpha$ be the integral group ring of Ψ_α . We view elements of Λ as non-commutative polynomials in the variables $\{a, a_\bullet\}_{a \in \alpha}$ subject to the relations $aa_\bullet = a_\bullet a$ and $a\tau(a) = a_\bullet\tau(a)_\bullet = 1$ for all $a \in \alpha$.

10.4. Generalized colorings. Let X be a left Λ -module. An X -coloring (with respect to $\beta \subset \alpha$) of a nanoword $(\mathcal{A}, w : \hat{n} \rightarrow \mathcal{A})$ is a function $f : \{0, 1, \dots, n\} \rightarrow X$ such that for any $A \in \mathcal{A}$ with $|A| = a \in \beta$,

$$f(i_A) = af(i_A - 1), \quad f(j_A) = a_\bullet f(j_A - 1) + (1 - aa_\bullet)f(i_A - 1)$$

and for any $A \in \mathcal{A}$ with $|A| = a \in \alpha - \beta$,

$$f(i_A) = a_\bullet f(i_A - 1) + (1 - aa_\bullet)f(j_A - 1), \quad f(j_A) = af(j_A - 1).$$

For $X = \mathbb{Z}/3\mathbb{Z}$, $ax = x$, $a_\bullet x = -x$ for all $a \in \alpha, x \in X$, this gives tricolorings of w .

The sum or the difference of two X -colorings of w is always an X -coloring of w . The zero function $f = 0$ is an X -coloring of w . Generally speaking, non-zero constant functions are not X -colorings unless we make additional assumptions like for instance $ax = x$ for all $a \in \alpha, x \in X$. If X is finite as a set, then there is only a finite number of X -colorings f of w with given input $k = f(0) \in X$ and output $l = f(n) \in X$. Denote this number $X(\beta, w)_{k,l}$. As in Sect. 10.1, either $X(\beta, w)_{k,l} = 0$ or $X(\beta, w)_{k,l} = X(\beta, w)_{0,0}$. An easy check shows that $X(\beta, w)_{k,l}$ is preserved under homotopy moves on w (we shall verify this in Sect. 11.3). The formula $w \mapsto (X(\beta, w)_{k,l})_{k,l \in X}$ defines a representation of $\mathcal{N}_\bullet(\alpha)$ in the monoid of integral $(\text{card}(X) \times \text{card}(X))$ -matrices with non-negative entries.

A vast family of Λ -modules can be obtained by the following construction. Pick a unital associative ring R and two functions $p, p_\bullet : \alpha \rightarrow R$ such that $p(a)p(\tau(a)) = p_\bullet(a)p_\bullet(\tau(a)) = 1$ for all $a \in \alpha$. Any left R -module X becomes a Λ -module by $ax = p(a)x$ and $a_\bullet x = p_\bullet(a)x$ for $a \in \alpha, x \in X$.

As an application, consider the nanoword $ABAB$ over $\alpha = \{a, \tau(a), b, \tau(b)\}$ with $|A| = a, |B| = b$. We check that this nanoword is not homotopic to its inverse. Set $X = R = \mathbb{Z}/5\mathbb{Z}$, $\beta = \{a, \tau(a)\}$, $p = 1$ (the constant function) and let $p_\bullet : \alpha \rightarrow \mathbb{Z}/5\mathbb{Z}$ be given by $p_\bullet(a) = p_\bullet(\tau(b)) = 2$ and $p_\bullet(\tau(a)) = p_\bullet(b) = 3$. The nanoword $ABAB$ admits a non-constant X -coloring

$$\begin{array}{ccccccc} & 0 & & 0 & & 3 & & 1 & & 1 \\ - & A & - & B & - & A & - & B & - \end{array}$$

while the inverse nanoword $A'B'A'B'$ with $|A'| = \tau(a), |B'| = \tau(b)$ admits only constant X -colorings.

10.5. Exercise. Prove the identity $X(\beta, \overline{w}^-)_{k,l} = X(\alpha - \beta, w)_{l,k}$ for any finite Λ -module X and $k, l \in X$.

11. MATRICES, MODULES, AND POLYNOMIALS

Throughout this section we fix a set $\beta \subset \alpha$ such that $\tau(\beta) = \beta$. We associate with each nanoword w a certain module $K_\beta(w)$. We begin with so-called marked modules and weighted matrices.

11.1. Marked modules and weighted matrices. By a *marked module* over a unital associative ring R , we mean a left R -module K endowed with an ordered pair of vectors $v_-, v_+ \in K$ called the *input* and the *output*, respectively. By *isomorphisms* of marked modules we mean isomorphisms of modules preserving the input and the output.

Let R^{ab} be the commutative ring obtained by quotienting R by the 2-sided ideal generated by the set $\{rs - sr \mid r, s \in R\}$. Denote the projection $R \rightarrow R^{ab}$ by q .

A *weighted matrix* over R is a finite matrix over R endowed with an ordered pair of elements of R^{ab} called the *weights*. Consider the following transformations on a weighted matrix M with $n \geq 1$ rows and $m \geq 1$ columns.

(i)' Permutation of two consecutive rows and simultaneous multiplication of both weights by -1 .

(ii)' Addition to the k -th column of the j -th column multiplied on the right by an element of R where $1 \leq k \leq m$, $2 \leq j \leq m-1$, and $k \neq j$. The weights are preserved.

(iii)' Striking out the i -th row and the j -th column provided $n \geq 2$, $1 \leq i \leq n$, $2 \leq j \leq m-1$ and the (i, j) -th entry $M_{i,j}$ of M is invertible in R and is the only non-zero entry in the i -th row. The first weight is multiplied by $(-1)^{i+j+1}q(M_{i,j})$ and the second weight is multiplied by $(-1)^{i+j}q(M_{i,j})$.

We say that two weighted matrices (possibly of different sizes) are *equivalent* if they can be obtained from each other by a finite sequence of transformations (i)' – (iii)' and the inverse transformations. The number $m - n$, called the *deficiency*

of M , is preserved under the transformations (i)' – (iii)'. For a weighted matrix M , denote by $\div M$ the weighted matrix obtained by multiplying both weights of M by -1 .

Each weighted $(n \times m)$ -matrix M presents an R -module denoted $K(M)$. More precisely, M determines an R -homomorphism $R^n \rightarrow R^m$ where the canonical basis vectors of R^m (resp. of R^n) correspond to the columns (resp. the rows) of M . The cokernel of this homomorphism is $K(M)$. The first and the last basis vectors of R^m project to certain $v_-, v_+ \in K(M)$. This gives a marked module $(K(M), v_-, v_+)$. It does not depend on the weights of M and is preserved (up to isomorphism) under the transformations (i)' – (iii)' on M and the involution $M \mapsto \div M$.

A weighted matrix $(M, r_- \in R^{ab}, r_+ \in R^{ab})$ of deficiency 1 gives rise to two elements of R^{ab} as follows. Consider the square matrix M_-^{ab} (resp. M_+^{ab}) obtained from M by deleting the first (resp. the last) column and projecting all entries to R^{ab} . For $\varepsilon \in \{+, -\}$, set $\nabla^\varepsilon(M) = r_\varepsilon \det(M_\varepsilon^{ab}) \in R^{ab}$. It is obvious that $\nabla^\varepsilon(M)$ is preserved under the transformations (i)' – (iii)' and $\nabla^\varepsilon(\div M) = -\nabla^\varepsilon(M)$. Note that M_ε^{ab} is a presentation matrix of the R^{ab} -module $R^{ab} \otimes_R K(M)/Rv_\varepsilon$. The principal ideal of R^{ab} generated by $\nabla^\varepsilon(M)$ coincides with the order ideal of this module. (This R^{ab} -module can be presented by a square matrix over R^{ab} and the determinant of any such matrix generates the order ideal.)

11.2. Matrices of nanowords. Recall the ring $\Lambda = \mathbb{Z}\Psi_\alpha$ introduced in Sect.

10.3. With each non-empty nanoword $(\mathcal{A}, w : \hat{n} \rightarrow \mathcal{A})$ over α we associate a $(2n \times (2n + 1))$ -matrix M over Λ as follows. Let us numerate all letters of $\mathcal{A}$ by the numbers $1, 2, \dots, n$. The k -th letter A_k determines the $(2k - 1)$ -st and $2k$ -th rows of M . To define these rows, we shall use notation of Sect. 8.1. If $a = |A_k| \in \beta$, then

$$M_{2k-1, i_A-1} = a, M_{2k-1, i_A} = -1, M_{2k, i_A-1} = 1 - aa_\bullet, M_{2k, j_A-1} = a_\bullet, M_{2k, j_A} = -1$$

while all other entries of these two rows are 0. If $a = |A| \in \alpha - \beta$, then one exchanges i_A, j_A in these formulas which gives

$$M_{2k-1, j_A-1} = a, M_{2k-1, j_A} = -1, M_{2k, i_A-1} = a_\bullet, M_{2k, i_A} = -1, M_{2k, j_A-1} = 1 - aa_\bullet$$

while all other entries of these two rows are 0. We provide M with weights as follows. The ring Λ^{ab} is the ring of polynomials in the commuting variables $\{a, a_\bullet\}_{a \in \alpha}$ subject to the relations $a\tau(a) = a_\bullet\tau(a)_\bullet = 1$ for $a \in \alpha$. For $a \in \alpha$, set $\langle a, w \rangle = \text{card}\{A \in \mathcal{A} \mid |A| = a\}$ and

$$r^w = \prod_{a \in \alpha - \beta} (-aa_\bullet)^{-\langle a, w \rangle} = \prod_{a \in \alpha - \beta} (-\tau(a)\tau(a)_\bullet)^{\langle a, w \rangle} \in \Lambda^{ab}.$$

The triple (M, r^w, r^w) is a weighted matrix over Λ denoted $M(w)$ or $M_\beta(w)$. Its equivalence class does not depend on the choice of numeration of letters of $\mathcal{A}$ since the permutation of the rows of M corresponding to two different choices is even. The next lemma describes the behavior of $M(w)$ under homotopy moves on w .

Lemma 11.2.1. *The equivalence class of $M(w)$ is preserved under the first and second homotopy moves on w . If a nanoword v is obtained from w by the third homotopy move, then $M(v)$ is equivalent to $\div M(w)$.*

Proof. It is obvious that isomorphic nanowords give the same weighted matrices. Consider the first homotopy move $w = (\mathcal{A}, xAAy) \mapsto (\mathcal{A} - \{A\}, xy) = v$ with $v \neq \emptyset$. We numerate the letters of $\mathcal{A}$ starting with A and set $a = |A| \in \alpha$. If $a \in \beta$, then

$$M(w) = \left(\begin{bmatrix} 0 & a & -1 & 0 & 0 \\ 0 & 1 - aa_{\bullet} & a_{\bullet} & -1 & 0 \\ m_1 & m_2 & 0 & m_3 & m_4 \end{bmatrix}, r^w, r^w \right)$$

where m_1, m_4 are matrices and m_2, m_3 are columns over Λ , all with the same number of rows equal to twice the length of v . It is clear that $r^v = r^w$ and $M(v) = ([m_1 | m_2 + m_3 | m_4], r^v, r^v)$ where the vertical bar separates submatrices. Multiplying the middle column of $M(w)$ by $a \in \Lambda$ and adding it to the previous column we obtain a matrix in which the first row contains only one non-zero entry -1 . Applying the transformation (iii)' and using that $aa_{\bullet} = a_{\bullet}a$, we obtain the weighted matrix

$$\left(\begin{bmatrix} 0 & 1 & -1 & 0 \\ m_1 & m_2 & m_3 & m_4 \end{bmatrix}, (-1)^{k+1}r^w, (-1)^k r^w \right)$$

where k is the number of columns in m_1 , that is the length of the word x . If $x \neq \emptyset$, then $k \neq 0$ and we can apply the transformation (ii)' adding the column containing m_2 to the next one. Applying then (iii)' to the first row we obtain $M(v)$. If $x = \emptyset$, then the assumption $v \neq \emptyset$ implies that $y \neq \emptyset$ so that the column containing m_3 is not the last one. We can apply the transformation (ii)' adding this column to the previous one. Then applying (iii)' to the first row we again obtain $M(v)$. If $a \in \alpha - \beta$, then

$$M(w) = \left(\begin{bmatrix} 0 & 0 & a & -1 & 0 \\ 0 & a_{\bullet} & -aa_{\bullet} & 0 & 0 \\ m_1 & m_2 & 0 & m_3 & m_4 \end{bmatrix}, r^w, r^w \right).$$

It is clear that $M(v) = ([m_1 | m_2 + m_3 | m_4], r^v, r^v)$ where $r^v = -aa_{\bullet}r^w$. Multiplying the middle column of $M(w)$ by $a^{-1} = \tau(a) \in \Lambda$ and adding it to the next column we obtain a matrix in which the first row contains only one non-zero entry a . Applying (iii)' to the first row we obtain the weighted matrix

$$\left(\begin{bmatrix} 0 & a_{\bullet} & -a_{\bullet} & 0 \\ m_1 & m_2 & m_3 & m_4 \end{bmatrix}, (-1)^k ar^w, (-1)^{k+1} ar^w \right)$$

where k is the length of x . If $x \neq \emptyset$, then we can add the column containing m_2 to the next one and then apply (iii)' to the first row to obtain $M(v)$. If $x = \emptyset$, then $y \neq \emptyset$ and we proceed as above.

Consider the move $w = (\mathcal{A}, xABYBAz) \mapsto (\mathcal{A} - \{A, B\}, xyz) = v$ where $|A| = \tau(|B|)$. Applying if necessary several inverse first homotopy moves to w , we can assume that the words x, y , and z are non-empty. We numerate the letters of $\mathcal{A}$ starting with A, B . Set $a = |A|$, $b = |B| = \tau(a)$, and $r = r^w \in \Lambda^{ab}$. By the

definition of Λ , we have $ab = a_\bullet b_\bullet = 1 \in \Lambda$. Note that $r^v = r^w = r$. This is obvious if $a, b \in \beta$ and follows from the equality $aa_\bullet bb_\bullet = aba_\bullet b_\bullet = 1$ in Λ^{ab} if $a, b \in \alpha - \beta$. Suppose that $a, b \in \beta$. Then

$$M(w) = \left(\begin{bmatrix} 0 & a & -1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 - aa_\bullet & 0 & 0 & 0 & 0 & a_\bullet & -1 & 0 \\ 0 & 0 & b & -1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 - bb_\bullet & 0 & 0 & b_\bullet & -1 & 0 & 0 \\ m_1 & m_2 & 0 & m_3 & m_4 & m_5 & 0 & m_6 & m_7 \end{bmatrix}, r, r \right)$$

where m_1, m_4, m_7 are non-void matrices and m_2, m_3, m_5, m_6 are columns over Λ , all with the same number of rows equal to twice the length of v . It is clear that

$$M(v) = ([m_1 \mid m_2 + m_3 \mid m_4 \mid m_5 + m_6 \mid m_7], r, r).$$

Multiplying by $b \in \Lambda$ the column of $M(w)$ containing m_3 and adding it to the previous column we obtain a matrix in which the third row contains only one non-zero entry -1 . Applying (iii)' to this row we obtain the weighted matrix

$$\left(\begin{bmatrix} 0 & a & -1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 - aa_\bullet & 0 & 0 & 0 & a_\bullet & -1 & 0 \\ 0 & 0 & 1 - bb_\bullet & 0 & b_\bullet & -1 & 0 & 0 \\ m_1 & m_2 & m_3 b & m_4 & m_5 & 0 & m_6 & m_7 \end{bmatrix}, (-1)^k r, (-1)^{k+1} r \right)$$

where k is the number of columns in m_1 , that is the length of the word x . Similarly using the entry -1 in the second row and applying (ii)', (iii)', we can transform the latter weighted matrix into

$$\begin{bmatrix} 0 & a & -1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 - bb_\bullet & 0 & b_\bullet & -1 & 0 \\ m_1 & m_2 + m_6(1 - aa_\bullet) & m_3 b & m_4 & m_5 & m_6 a_\bullet & m_7 \end{bmatrix}$$

with both weights equal to $(-1)^{l+1} r$ where l is the number of columns in m_4 . The latter matrix is similarly transformed into

$$\begin{bmatrix} 0 & (1 - bb_\bullet)a & 0 & b_\bullet & -1 & 0 \\ m_1 & m_2 + m_6(1 - aa_\bullet) + m_3 ba & m_4 & m_5 & m_6 a_\bullet & m_7 \end{bmatrix}$$

with weights $(-1)^{k+l} r, (-1)^{k+l+1} r$ and finally into

$$([m_1 \mid m_2 + m_6(1 - aa_\bullet) + m_3 ba + m_6 a_\bullet(1 - bb_\bullet)a \mid m_4 \mid m_5 + m_6 a_\bullet b_\bullet \mid m_7], r, r).$$

Since $ab = b_\bullet a_\bullet = 1$, the latter weighted matrix coincides with $M(v)$.

The case where $a, b \in \alpha - \beta$ is treated similarly. For completeness, we specify the matrix $M(w)$ in this case leaving to the reader to verify that it is equivalent to $M(v)$:

$$M(w) = \left(\begin{bmatrix} 0 & 0 & 0 & 0 & 0 & 0 & a & -1 & 0 \\ 0 & a_\bullet & -1 & 0 & 0 & 0 & 1 - aa_\bullet & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & b & -1 & 0 & 0 \\ 0 & 0 & b_\bullet & -1 & 0 & 1 - bb_\bullet & 0 & 0 & 0 \\ m_1 & m_2 & 0 & m_3 & m_4 & m_5 & 0 & m_6 & m_7 \end{bmatrix}, r, r \right).$$

Consider the move $w = (\mathcal{A}, xAByACzBCt) \mapsto (\mathcal{A}, xBAyCAzCBt) = v$ where $|A| = |B| = |C|$. Applying if necessary several inverse first homotopy moves to w , we can assume that the words x, y, z, t are non-empty. Set $a = |A| \in \alpha$, $q = 1 - aa_\bullet \in \Lambda$, and $r = r^w \in \Lambda^{ab}$. Clearly, $r^v = r^w = r$. If $a \in \beta$, then

$$M(w) = \begin{bmatrix} 0 & a & -1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & q & 0 & 0 & 0 & a_\bullet & -1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & a & -1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & q & 0 & 0 & 0 & 0 & 0 & 0 & a_\bullet & -1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & a & -1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & q & 0 & 0 & 0 & a_\bullet & -1 & 0 \\ m_1 & m_2 & 0 & m_3 & m_4 & m_5 & 0 & m_6 & m_7 & m_8 & 0 & m_9 & m_{10} \end{bmatrix}$$

with weights r, r and

$$M(v) = \begin{bmatrix} 0 & a & -1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & q & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & a_\bullet & -1 & 0 \\ 0 & 0 & a & -1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & q & 0 & 0 & 0 & a_\bullet & -1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & a & -1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & q & 0 & 0 & 0 & a_\bullet & -1 & 0 & 0 \\ m_1 & m_2 & 0 & m_3 & m_4 & m_5 & 0 & m_6 & m_7 & m_8 & 0 & m_9 & m_{10} \end{bmatrix}$$

with weights r, r . It is an exercise in linear algebra to verify that $M(w)$ is equivalent to the matrix

$$[m_1 \mid m_2 + m_3a^2 + m_6aq + m_9q \mid m_4 \mid m_5 + m_6aa_\bullet + m_9a_\bullet q \mid m_7 \mid m_8 + m_9a_\bullet^2 \mid m_{10}]$$

with weights r, r . The matrix $M(v)$ is equivalent to the same matrix with weights $-r, -r$. Thus, $M(v)$ is equivalent to $\div M(w)$. Similarly, if $a \in \alpha - \beta$, then $M(w)$ is equivalent to the matrix

$$[m_1 \mid m_2 + m_3a_\bullet^2 \mid m_4 \mid m_5 + m_3a_\bullet q + m_6aa_\bullet \mid m_7 \mid m_8 + m_3q + m_6aq + m_9a^2 \mid m_{10}]$$

with weights $-r, -r$ and $M(v)$ is equivalent to the same matrix with weights r, r . $\square$

To describe the behavior of the weighted matrix under operations on nanowords, we need two involutions on the set of weighted matrices over Λ . Define first a ring involution $\lambda \mapsto \bar{\lambda} : \Lambda \rightarrow \Lambda$ by $\bar{a} = a^{-1} = \tau(a)$ and $\overline{a_\bullet} = a_\bullet^{-1} = \tau(a)_\bullet$ for $a \in \alpha$. Given a weighted matrix M , denote by $\overline{M}$ the matrix obtained by applying this ring involution to all entries of M and applying the induced involution of Λ^{ab} to the weights. For an $(n \times m)$ -matrix $M = (M_{i,j})_{i,j}$, set $M^- = (M_{n+1-i, m+1-j})_{i,j}$. The second involution on the set of weighted matrices over Λ transforms (M, r_-, r_+) into (M^-, r_+, r_-) if the deficiency of M is odd and into (M^-, r_-, r_+) if the deficiency of M is even. Both involutions are compatible with the equivalence, that is transform equivalent weighted matrices into equivalent weighted matrices.

It follows from the definitions that $M(\overline{w}) = \overline{M(w)}$. The next lemma computes $M_{\alpha-\beta}(w^-)$ from $M_\beta(w)$.

Lemma 11.2.2. *For any nanoword $(\mathcal{A}, w)$, the weighted matrix $M_{\alpha-\beta}(w^-)$ is obtained from $\overline{M_\beta(w)}^-$ by transformations (i)' – (iii)' and the following transformations on weighted matrices: (iv)' addition to a row of another row multiplied on the left by an element of Λ , the weights being preserved and (v)' multiplication of a row by an invertible element $\lambda \in \Lambda$, the weights being multiplied by λ^{-1} .*

Proof. Let us numerate the letters of $\mathcal{A}$. Recall that the k -th letter A_k contributes 2 rows to $M_\beta(w)$. Set $a = |A_k|$, $i = i_{A_k}$, $j = j_{A_k}$. Assume first that $i \leq j - 2$. If $a \in \beta$, then the rows contributed by A_k are

$$\begin{bmatrix} 0_{i-1} & a & -1 & 0_{j-i-2} & 0 & 0 & 0_{n-j} \\ 0_{i-1} & 1 - aa_\bullet & 0 & 0_{j-i-2} & a_\bullet & -1 & 0_{n-j} \end{bmatrix}$$

where 0_s stands for a row of s zeros. The corresponding rows in $\overline{M_\beta(w)}^-$ are

$$\begin{bmatrix} 0_{n-j} & -1 & a_\bullet^{-1} & 0_{j-i-2} & 0 & 1 - a^{-1}a_\bullet^{-1} & 0_{i-1} \\ 0_{n-j} & 0 & 0 & 0_{j-i-2} & -1 & a^{-1} & 0_{i-1} \end{bmatrix}.$$

Permuting these two rows we obtain

$$\begin{bmatrix} 0_{n-j} & 0 & 0 & 0_{j-i-2} & -1 & a^{-1} & 0_{i-1} \\ 0_{n-j} & -1 & a_\bullet^{-1} & 0_{j-i-2} & 0 & 1 - a^{-1}a_\bullet^{-1} & 0_{i-1} \end{bmatrix}.$$

Multiply the first row by $a_\bullet^{-1} - a$ and add it to the second row. This gives

$$\begin{bmatrix} 0_{n-j} & 0 & 0 & 0_{j-i-2} & -1 & a^{-1} & 0_{i-1} \\ 0_{n-j} & -1 & a_\bullet^{-1} & 0_{j-i-2} & a - a_\bullet^{-1} & 0 & 0_{i-1} \end{bmatrix}.$$

Multiplying the first row by $-a$ and the second row by $-a_\bullet$ we obtain

$$\begin{bmatrix} 0_{n-j} & 0 & 0 & 0_{j-i-2} & a & -1 & 0_{i-1} \\ 0_{n-j} & a_\bullet & -1 & 0_{j-i-2} & 1 - aa_\bullet & 0 & 0_{i-1} \end{bmatrix}.$$

These are exactly the rows contributed by A_k to $M_{\alpha-\beta}(w^-)$. The cases where $i = j - 1$ or $a \in \alpha - \beta$ are similar. That the weights behave appropriately is left to the reader as an exercise. $\square$

Given two nanowords w, w' , we can compute the weighted matrix $M(ww')$ from $M(w) = (M, r_-, r_+)$ and $M(w') = (M', r'_-, r'_+)$. Let us write $M = [M_+ \ m_+]$ where m_+ is the last column of M and M_+ is obtained from M by deleting m_+ . Similarly, $M' = [m'_- \ M'_-]$ where m'_- is the first column of M' . Then

$$M(ww') = \left(\begin{bmatrix} M_+ & m_+ & 0 \\ 0 & m'_- & M'_- \end{bmatrix}, r_- r'_-, r_+ r'_+ \right).$$

11.3. Module $K_\beta(w)$. By Sect. 11.1, the weighted matrix $M_\beta(w)$ of a nanoword w gives rise to a marked Λ -module $K_\beta(w) = K(M_\beta(w))$. Lemma 11.2.1 implies that $K_\beta(w)$ is preserved under homotopy moves on w . It is clear that $K_\beta(\overline{w}) = \overline{K_\beta(w)}$ where for a Λ -module K , the Λ -module $\overline{K}$ is obtained from K by pulling back along the involution $\lambda \mapsto \overline{\lambda} : \Lambda \rightarrow \Lambda$. Lemma 11.2.2 implies that $K_{\alpha-\beta}(w^-)$ is obtained from $\overline{K_\beta(w)}$ by permuting the input with the output.

For a Λ -module X , the X -colorings of w bijectively correspond to Λ -linear homomorphisms $K_\beta(w) \rightarrow X$. Hence for finite (as a set) X , the number of X -colorings of w with given input and output is a homotopy invariant of w .

Further homotopy invariants of w are provided by the elementary ideals and other invariants of the Λ^{ab} -module $\Lambda^{ab} \otimes_\Lambda K_\beta(w)$ and of its quotients by the input and/or the output. We discuss two such invariants in the next subsection.

11.4. Polynomials $\nabla_\beta^\pm(w)$. Since the deficiency of the weighted matrix $M = M_\beta(w)$ is equal to 1, Sect. 11.1 yields two polynomials $\nabla^-(M), \nabla^+(M) \in \Lambda^{ab}$. Lemma 11.2.1 implies that they are homotopy invariants of w at least up to multiplication by -1 . We eliminate the sign indeterminacy as follows. Consider the homomorphism $\text{aug} : \Lambda^{ab} \rightarrow \mathbb{Z}$ sending all $a, a_\bullet$ to 1. In other words, the value of aug on $\lambda \in \Lambda^{ab}$ is the sum of coefficients of λ . Applying aug to all entries of M and permuting the rows, we obtain the matrix

$$\begin{bmatrix} 1 & -1 & 0 & 0 & \cdots \\ 0 & 1 & -1 & 0 & \cdots \\ & & \cdots & & \\ \cdots & 0 & 0 & 1 & -1 \end{bmatrix}.$$

Therefore $\text{aug}(\nabla^\varepsilon(M)) = \pm 1$ for $\varepsilon = \pm 1$. The polynomial

$$\nabla_\beta^\varepsilon(w) = \text{aug}(\nabla^\varepsilon(M)) \nabla^\varepsilon(M) \in \Lambda^{ab}$$

is a homotopy invariant of w without any indeterminacy. Clearly, $\text{aug}(\nabla_\beta^\varepsilon(w)) = 1$. The results of Sect. 11.2 imply that

$$\nabla_\beta^\varepsilon(\overline{w}) = \overline{\nabla_\beta^\varepsilon(w)}, \quad \nabla_{\alpha-\beta}^\varepsilon(w^-) = \overline{\nabla_\beta^{-\varepsilon}(w)}, \quad \nabla_\beta^\varepsilon(ww') = \nabla_\beta^\varepsilon(w) \nabla_\beta^\varepsilon(w').$$


12. INVARIANT λ

We use the modules of nanowords to define a Λ -valued homotopy invariant of nanowords λ . As an application we give a homotopy classification of nanowords of length 6.

12.1. Invariants λ' and λ . For a nanoword $(\mathcal{A}, w : \hat{n} \rightarrow \mathcal{A})$, consider the Λ -module $K_\alpha(w)$ corresponding to $\beta = \alpha$. This module is free of rank 1. Indeed, let $v_- = x_0, x_1, \dots, x_n = v_+$ be the generators of $K_\alpha(w)$ determined by the first, second, etc. columns of the matrix $M = M_\alpha(w)$. The two rows of M derived from a letter $A \in \mathcal{A}$ express x_{i_A} as a linear combination of x_s with $s < i_A$ and express x_{j_A} as a linear combination of x_s with $s < j_A$. Therefore $K_\alpha(w) = \Lambda v_-$. Then $v_+ = \lambda'(w)v_-$ for a unique $\lambda'(w) \in \Lambda$ which is a homotopy invariant of w . It admits an equivalent but more convenient version $\lambda(w)$ defined as follows. Consider the involutive anti-automorphism $\iota : \Lambda \rightarrow \Lambda$ equal to the identity on the generators $\{a, a_\bullet\}_{a \in \alpha} \subset \Lambda$. Thus ι acts on monomials by reading them from right to left. For instance, $\iota(aba_\bullet) = a_\bullet ba$. Set

$$\lambda(w) = \iota(\lambda'(w)) \in \Lambda.$$

We describe a simple method allowing to compute $\lambda(w)$ (and extending the Silver-Williams method [SW] in the setting of virtual strings). For each $i = 1, \dots, n$


 FIGURE 3. Graph Γ_w for $w = ABAB$

there is a unique row in M expressing the generator $x_i \in K_\alpha(w)$ via the generators with smaller indices. This relation has either the form $x_i = k_i x_{i-1}$ with $k_i \in \Lambda$ or the form $x_i = k_i x_{i-1} + l_i x_{i'}$ with $k_i, l_i \in \Lambda$, $l_i \neq 0$, and $i' < i - 1$. We form a graph Γ_w with $n + 1$ vertices $\{0, 1, 2, \dots, n\}$ and $3n/2$ edges as follows. A relation of the form $x_i = k_i x_{i-1}$ gives rise to one edge - the segment $[i - 1, i] \subset \mathbf{R}$ with endpoints $i - 1, i$. We place on this edge the weight k_i . A relation of the form $x_i = k_i x_{i-1} + l_i x_{i'}$ gives rise to two edges: the segment $[i - 1, i] \subset \mathbf{R}$ and an arc connecting i with i' (and disjoint from the rest of Γ_w). We place on these two edges the weights k_i and l_i , respectively. Consider a path leading from n to 0 and following along the edges of Γ_w , always from the bigger endpoint to the smaller one. Compute the product of weights associated with the edges lying on the path writing the weights from right to left in the order of appearance of the edges on the path. Summing these products over all such paths from n to 0 we obtain $\lambda(w)$.

Example: for $w = ABAB$ with $|A| = a \in \alpha$, $|B| = b \in \alpha$, there are three paths in Γ_w contributing to $\lambda(w)$, cf. Figure 3. This gives

$$(12.1.1) \quad \lambda(w) = aba_\bullet b_\bullet + (1 - aa_\bullet)b_\bullet + a(1 - bb_\bullet).$$

We describe a more general class of nanowords for which λ can be computed explicitly. Pick a finite sequence of letters (possibly with repetitions) $a_1, \dots, a_m \in \alpha$ and a permutation $\sigma : \hat{m} \rightarrow \hat{m}$. Consider the nanoword $(\mathcal{A} = \{A_1, \dots, A_m\}, w : \widehat{2m} \rightarrow \mathcal{A})$ where $|A_i| = a_i$, $w(i) = A_i$ and $w(i + m) = A_{\sigma(i)}$ for $i = 1, \dots, m$. Thus $w = A_1 A_2 \cdots A_m A_{\sigma(1)} A_{\sigma(2)} \cdots A_{\sigma(m)}$ with $|A_i| = a_i$. The corresponding graph Γ contains precisely $m + 1$ paths from $2m$ to 0 and

$$(12.1.2) \quad \lambda(w) = a_1 a_2 \cdots a_m (a_{\sigma(1)})_\bullet (a_{\sigma(2)})_\bullet \cdots (a_{\sigma(m)})_\bullet \\ + \sum_{i=1}^m a_1 a_2 \cdots a_{\sigma(i)-1} (1 - a_{\sigma(i)} (a_{\sigma(i)})_\bullet) (a_{\sigma(i+1)})_\bullet (a_{\sigma(i+2)})_\bullet \cdots (a_{\sigma(m)})_\bullet.$$

12.2. Properties of λ . It follows from the definitions that $\lambda(\overline{w}) = \overline{\lambda(w)}$ and $\lambda(ww') = \lambda(w)\lambda(w')$ for any nanowords w, w' . The graph Γ_{w^-} is obtained from Γ_w by relabeling the vertices (i becomes $n - i$) and exchanging $a \leftrightarrow a_\bullet$ for all $a \in \alpha$. Therefore $\lambda(w^-) = \kappa(\lambda(w))$ where κ is the ring anti-automorphism of Λ defined

on the generators $a \in \alpha$ by $\kappa(a) = a_\bullet, \kappa(a_\bullet) = a$. Thus if w is homotopically symmetric then $\kappa(\lambda(w)) = \lambda(w)$. If w is homotopically skew-symmetric then $\kappa(\lambda(w)) = \lambda(w)$. If w is contractible, then $\lambda(w) = 1$.

To state further properties of $\lambda(w)$, recall the group Π defined in Sect. 4.1 and consider the ring homomorphisms $p, r, r_\bullet : \Lambda \rightarrow \mathbb{Z}\Pi$ defined on the generators of Λ by $p(a) = z_a, p(a_\bullet) = z_{\tau(a)}, r(a) = z_a, r(a_\bullet) = 1$ and $r_\bullet(a) = 1, r_\bullet(a_\bullet) = z_a$.

Lemma 12.2.1. *For any nanoword w , $p(\lambda(w)) = \gamma(w) \in \Pi$ is the invariant defined in Sect. 4.1 and $r(\lambda(w)) = r_\bullet(\lambda(w)) = 1 \in \mathbb{Z}\Pi$.*

Proof. Substituting $a_\bullet = a^{-1}$ in the matrix $M_\alpha(w)$ we obtain $x_{i_A+1} = |A|x_{i_A}$ and $x_{j_A+1} = |A|^{-1}x_{j_A}$. Comparing with the definition of $\gamma(w)$, we obtain $p(\lambda(w)) = \gamma(w)$. Substituting $a_\bullet = 1$ in the matrix $M_\alpha(w)$ and adding all rows we obtain a row with first entry 1, last entry -1 , and all other entries 0. Hence $r(\lambda(w)) = 1$. Substituting $\lambda(w^-) = \kappa(\lambda(w))$ in $r(\lambda(w^-)) = 1$ we obtain a formula equivalent to $r_\bullet(\lambda(w)) = 1$. $\square$

The polynomials $\nabla_\alpha^-(w), \nabla_\alpha^+(w)$ defined in Sect. 11.4 (for $\beta = \alpha$) can be computed as follows: $\nabla_\alpha^-(w) = 1$ and $\nabla_\alpha^+(w) = q(\lambda'(w)) = q(\lambda(w))$ where q is the projection $\Lambda \rightarrow \Lambda^{ab}$. Thus $\lambda(w)$ is a common extension of $\nabla_\alpha^+(w)$ and $\gamma(w)$.

12.3. Derived invariants. For a monomial $m \in \Lambda$ in the generators $\{a, a_\bullet\}_{a \in \alpha}$, denote $\deg(m)$ the number of generators without $\bullet$ appearing in m and denote $\deg_\bullet(m)$ the number of generators with $\bullet$ in m . For instance $\deg(aba_\bullet) = 2$ and $\deg_\bullet(aba_\bullet) = 1$. We have

$$\Lambda = \Lambda_{0,0} \oplus \Lambda_{0,1} \oplus \Lambda_{1,0} \oplus \Lambda_{1,1}$$

where $\Lambda_{i,j} \subset \Lambda$ is additively generated by the monomials m such that $\deg(m) = i \pmod{2}$ and $\deg_\bullet(m) = j \pmod{2}$. As a corollary, the invariant $\lambda(w)$ of a nanoword w splits as a sum of 4 homotopy invariants

$$\lambda(w) = \lambda_{0,0}(w) + \lambda_{0,1}(w) + \lambda_{1,0}(w) + \lambda_{1,1}(w)$$

such that $\lambda_{i,j}(w) \in \Lambda_{i,j}$ for all i, j .


Consider the ring homomorphism $\psi : \Lambda \rightarrow \mathbb{Z}\Pi \otimes_{\mathbb{Z}} \mathbb{Z}\Pi$ defined on the generators $a \in \alpha$ of Λ by $\psi(a) = z_a \otimes 1$ and $\psi(a_\bullet) = 1 \otimes z_a$. For instance, $\psi(aba_\bullet) = z_a z_b \otimes z_a$. The set $\{x \otimes y\}_{x,y \in \Pi}$ is a basis of the additive group $\mathbb{Z}\Pi \otimes_{\mathbb{Z}} \mathbb{Z}\Pi$. For $\lambda \in \Lambda$, we have a unique (finite) expansion $\psi(\lambda) = \sum_{x,y \in \Pi} \lambda^{x,y} x \otimes y$ with $\lambda^{x,y} \in \mathbb{Z}$. For a nanoword w , this gives a set of homotopy invariants $\{(\lambda(w))^{x,y} \in \mathbb{Z}\}_{x,y \in \Pi}$.

12.4. Proof of Lemma 9.4.1. For $w = A_3 A_2 A_3 A_1 B A_2 A_1 B$, there are 7 paths in the graph Γ_w contributing to $\lambda(w)$, see Figure 4. We have

$$\begin{aligned} \lambda(A_3 A_2 A_3 A_1 B A_2 A_1 B) &= (1 - aa_\bullet)^2 b_\bullet + (1 - aa_\bullet)a(1 - bb_\bullet) + (1 - aa_\bullet)aba_\bullet^2 b_\bullet \\ &\quad + a(1 - aa_\bullet)a_\bullet b_\bullet + a^2 a_\bullet(1 - aa_\bullet)b_\bullet + a^3 a_\bullet(1 - bb_\bullet) + a^3 a_\bullet ba_\bullet^2 b_\bullet. \end{aligned}$$

This implies that $\lambda_{1,1}(A_3 A_2 A_3 A_1 B A_2 A_1 B) = a^3 a_\bullet - a^3 a_\bullet^2 b_\bullet$. In particular if $a = \tau(a)$, then we have $\lambda_{1,1}(A_3 A_2 A_3 A_1 B A_2 A_1 B) = aa_\bullet - ab_\bullet$. Similarly,

$$\lambda(A_3 A_2 B A_3 A_1 A_2 A_1 B) = (1 - aa_\bullet)^2 b_\bullet + (1 - aa_\bullet)aa_\bullet^2 b_\bullet + a(1 - aa_\bullet)a_\bullet b_\bullet$$


 FIGURE 4. Graph Γ_w for $w = A_3A_2A_3A_1BA_2A_1B$

$$+a^2(1-bb_\bullet) + a^2ba_\bullet(1-aa_\bullet)b_\bullet + a^2baa_\bullet^3b_\bullet.$$

This implies that $\lambda_{1,1}(A_3A_2BA_3A_1A_2A_1B) = aa_\bullet^2b_\bullet - bb_\bullet$. For $a = \tau(a)$, we have $\lambda_{1,1}(A_3A_2BA_3A_1A_2A_1B) = ab_\bullet - bb_\bullet$. The equality $aa_\bullet - ab_\bullet = ab_\bullet - bb_\bullet$ implies that $aa_\bullet = ab_\bullet$ so that $a = b$ which contradicts the assumptions of the lemma.

12.5. Proof of Lemma 9.7.1. A desingularization of $w_{a,b}^6$ gives the nanoword $A_3A_2BA_3A_1BA_2A_1$ in the α -alphabet (9.2.1). We have

$$\begin{aligned} \lambda(A_3A_2BA_3A_1BA_2A_1) &= (1-aa_\bullet)^2 + (1-aa_\bullet)ab_\bullet a_\bullet^2 + a(1-aa_\bullet)a_\bullet \\ &\quad + a^2(1-bb_\bullet)a_\bullet^2 + a^2ba_\bullet(1-aa_\bullet) + a^2baa_\bullet b_\bullet a_\bullet^2. \end{aligned}$$

Substituting here $a^2 = a_\bullet^2 = b^2 = b_\bullet^2 = 1$ (due to the assumptions $a = \tau(a), b = \tau(b)$), we easily obtain that

$$\lambda_{1,1} = \lambda_{1,1}(A_3A_2BA_3A_1BA_2A_1) = 2 - ba - a_\bullet b_\bullet + baa_\bullet b_\bullet.$$

Note that $ba \neq 1$ and $a_\bullet b_\bullet \neq 1$ since $\tau(a) = a \neq b$. Therefore $\lambda_{1,1} \neq 0$. Moreover, knowing $\lambda_{1,1}$ we can recover ba : This is the only non-trivial element $x \in \Pi$ such that $(\lambda_{1,1})^{x,1} = -1$. The equality $ba = b'a'$ implies that $b = b', a = a'$ which concludes the proof of the lemma.

12.6. Remark. For a nanoword $w = A_1 \cdots A_n$, set $w_* = |A_1|_* \cdots |A_n|_* \in \Lambda$ where $|A_i|_* = |A_i|$ if i numerates the first occurrence of the letter A_i in w and $|A_i|_* = |A_i|_\bullet$ if i numerates the second occurrence of A_i in w . The computation of λ in Sect. 12.1 implies that $\lambda(w) - w_*$ is an algebraic sum of monomials m in the generators $a, a_\bullet$ such that $\max(\deg(m), \deg_\bullet(m)) \leq n/2$ and $\deg(m) + \deg_\bullet(m) < n$.

12.7. Exercise. Deduce from Lemma 12.2.1 that if α is a one-element set, then $\lambda(w) = 1$ for any nanoword w over α .

13. ANALYSIS LITTERAE II

In this section we give a homotopy classification of nanowords of length ≤ 6 . We begin with nanowords of length 4. We shall use the homomorphism ψ introduced in Sect. 12.3.

13.1. Nanowords of length 4 re-examined. Let $w = ABAB$ with $|A| = a \in \alpha, |B| = b \in \alpha$. Formula 12.1.1 implies that

$$\lambda_{0,0}(w) = aba \bullet b \bullet, \lambda_{1,0}(w) = a - aa \bullet b \bullet, \lambda_{0,1}(w) = b \bullet - abb \bullet, \lambda_{1,1}(w) = 0.$$

Then $\psi(\lambda_{0,1}(w)) = (1 - z_a z_b) \otimes z_b$. Note that $z_a z_b = 1 \in \Pi$ if and only if $a = \tau(b)$. Therefore $\psi(\lambda_{0,1}(w)) = 0$ implies that $a = \tau(b)$ and then w is contractible. Conversely, if w is contractible then $\lambda_{0,1}(w) = 0$. We conclude that w is contractible if and only if $\psi(\lambda_{0,1}(w)) = 0$. If $\psi(\lambda_{0,1}(w)) \neq 0$, then we can recover a and b from $\psi(\lambda_{0,1}(w))$. This gives an alternative proof of Theorem 8.4.1 and shows that λ is a faithful homotopy invariant of nanowords of length 4.

13.2. Nanowords of length 6. Pick three letters $a, b, c \in \alpha$ (possibly coinciding). Let $\mathcal{A}$ be the α -alphabet consisting of 3 letters A, B, C with $|A| = a, |B| = b, |C| = c$. Consider the nanowords $w_{a,b,c}^1 = ABCABC$ and

$$w_{a,b,c}^2 = ABCACB, w_{a,b,c}^3 = ABCBAC, w_{a,b,c}^4 = ABCBCA, w_{a,b,c}^5 = ABACBC$$

in this α -alphabet. It is easy to check that any nanoword of length 6 is either homotopic to a nanoword of length ≤ 4 or is isomorphic to $w_{a,b,c}^i$ with $i \in \{1, \dots, 5\}$. We now point out obvious sufficient conditions for $w_{a,b,c}^i$ to be contractible.

If $a = \tau(b)$ or $c = \tau(b)$, then $w_{a,b,c}^1 = ABCABC$ is contractible. We say that an (ordered) triple $a, b, c \in \alpha$ is *1-regular* if $a \neq \tau(b) \neq c$.

If $c = \tau(b)$, then $w_{a,b,c}^2 = ABCACB$ is contractible. A triple $a, b, c \in \alpha$ is *2-regular* if $c \neq \tau(b)$.

If $a = \tau(b)$, then $w_{a,b,c}^3 = ABCBAC$ is contractible. A triple $a, b, c \in \alpha$ is *3-regular* if $a \neq \tau(b)$.

If $c = \tau(b)$, then $w_{a,b,c}^4 = ABCBCA$ is contractible. A triple $a, b, c \in \alpha$ is *4-regular* if $c \neq \tau(b)$. (This coincides with the 2-regularity).

If $a = b = c = \tau(a)$, then $w_{a,b,c}^5 = ABACBC$ is contractible by the moves $ABACBC \mapsto BACACB \mapsto BB \mapsto \emptyset$. We say that a triple $a, b, c \in \alpha$ is *singular* if $a = b = c = \tau(a)$ and *5-regular* otherwise.

The following theorem gives a homotopy classification of nanowords of length 6. Its proof shows that λ is a faithful homotopy invariant of nanowords of length 6 corresponding to regular triples a, b, c . However, one of the claims of the theorem - that $w_{a,b,c}^5$ is not homotopic to a nanoword of length 4 - needs subtler techniques in the case $a = c = \tau(b) \neq b$. This is due to the fact that in this case the invariant λ and all the other invariants of nanowords introduced above do not distinguish $w_{a,b,c}^5 = ABACBC$ from $w_{a,c} = ACAC$. That these nanowords are not homotopic will be proven in Sect. 16.5.

Theorem 13.2.1. *For $i = 1, \dots, 5$ and any i -regular triple $a, b, c \in \alpha$, the nanoword $w_{a,b,c}^i$ is neither contractible nor homotopic to a nanoword of length 4. The nanowords w^i corresponding to i -regular triples a, b, c and a', b', c' are homotopic if and only if $a = a', b = b',$ and $c = c'$. For $i \neq j$, the nanowords w^i corresponding to i -regular triples are not homotopic to nanowords w^j corresponding to j -regular triples with one exception: $w_{a,b,c}^4 \simeq w_{a,b,c}^5$ for $a = b = c \neq \tau(a)$.*

Proof. Set $w = w_{a,b,c}^1$. Formula (12.1.2) with $\sigma = \text{id}$ yields

$$\lambda(w) = abca_{\bullet}b_{\bullet}c_{\bullet} + (1 - aa_{\bullet})b_{\bullet}c_{\bullet} + a(1 - bb_{\bullet})c_{\bullet} + ab(1 - cc_{\bullet}).$$

Therefore $\lambda_{0,1}(w) = \lambda_{1,0}(w) = 0$ and

$$\lambda_{0,0}(w) = ab + b_{\bullet}c_{\bullet} - abb_{\bullet}c_{\bullet}, \quad \lambda_{1,1}(w) = ac_{\bullet} - aa_{\bullet}b_{\bullet}c_{\bullet} - abcc_{\bullet} + abca_{\bullet}b_{\bullet}c_{\bullet}.$$

Clearly, $\psi(1 - \lambda_{0,0}(w)) = (1 - z_a z_b) \otimes (1 - z_b z_c)$. If w is contractible, then $\lambda_{0,0}(w) = 1$ and we must have $z_a z_b = 1$ or $z_b z_c = 1$. Then $a = \tau(b)$ or $c = \tau(b)$ which is excluded by the 1-regularity.

This argument shows, moreover, that we can recover $z_a z_b$ and $z_b z_c$ from $\lambda_{0,0}(w)$. Therefore if $w_{a,b,c}^1 \simeq w_{a',b',c'}^1$, then $z_a z_b = z_{a'} z_{b'}$ and $z_b z_c = z_{b'} z_{c'}$. The equality $z_a z_b = z_{a'} z_{b'} \in \Pi$ holds in only two cases: when $a = a', b = b'$ and when $a = \tau(b)$ and $a' = \tau(b')$. The second option is excluded by the 1-regularity. Therefore $a = a', b = b',$ and $c = c'$.

That w is not homotopic to a nanoword of length 4 follows from the formula $\lambda_{0,1}(w) = 0$ and the fact that nanowords of length 4 with $\lambda_{0,1} = 0$ are contractible.

Set $w = w_{a,b,c}^2$. Formula (12.1.2) with $\sigma(1) = 1, \sigma(2) = 3, \sigma(3) = 2$ yields

$$\lambda(w) = abca_{\bullet}c_{\bullet}b_{\bullet} + (1 - aa_{\bullet})c_{\bullet}b_{\bullet} + ab(1 - cc_{\bullet})b_{\bullet} + a(1 - bb_{\bullet}).$$

Therefore $\lambda_{0,0}(w) = c_{\bullet}b_{\bullet}, \lambda_{0,1}(w) = 0$, and

$$\lambda_{1,0}(w) = a - abcc_{\bullet}b_{\bullet}, \quad \lambda_{1,1}(w) = abca_{\bullet}c_{\bullet}b_{\bullet} - aa_{\bullet}c_{\bullet}b_{\bullet}.$$

If w is contractible, then $\lambda_{1,0}(w) = 0$. This implies $c = \tau(b)$, excluded by the 2-regularity. For a 2-regular triple a, b, c , we can recover b, c from $\lambda_{0,0}(w)$ and a from $\lambda_{1,0}(w)$. (Indeed, $x = z_a$ is the only element of Π such that $(\lambda_{1,0}(w))^{x,1} \neq 0$.) That w is not homotopic to a nanoword of length 4 follows from the formula $\lambda_{0,1}(w) = 0$ and the fact that a nanoword of length 4 with $\lambda_{0,1} = 0$ must be contractible. That w is not homotopic to a nanoword of type w^1 follows from the formulas $\lambda_{1,0}(w^1) = 0$ and $\lambda_{1,0}(w) \neq 0$.

Set $w = w_{a,b,c}^3$. Formula (12.1.2) with $\sigma(1) = 2, \sigma(2) = 1, \sigma(3) = 3$ yields

$$\lambda(w) = abcb_{\bullet}a_{\bullet}c_{\bullet} + a(1 - bb_{\bullet})a_{\bullet}c_{\bullet} + (1 - aa_{\bullet})c_{\bullet} + ab(1 - cc_{\bullet}).$$

Therefore $\lambda_{0,0}(w) = ab, \lambda_{1,0}(w) = 0$, and

$$\lambda_{0,1}(w) = c_{\bullet} - abb_{\bullet}a_{\bullet}c_{\bullet}, \quad \lambda_{1,1}(w) = abcb_{\bullet}a_{\bullet}c_{\bullet} - abcc_{\bullet}.$$

If w is contractible, then $\lambda_{0,1}(w) = 0$. This implies $a = \tau(b)$, excluded by the 3-regularity. For a 3-regular triple a, b, c , we can recover a, b from $\lambda_{0,0}(w)$ and c from $\lambda_{0,1}(w)$. (Indeed, $y = z_c$ is the only element of Π such that $(\lambda_{0,1}(w))^{1,y} \neq 0$.) That w is not homotopic to a nanoword of length 4 follows from the formula $\lambda_{1,0}(w) = 0$ and the fact that a nanoword of length 4 with $\lambda_{1,0} = 0$ must be contractible. That w is not homotopic to a nanoword of type w^1 or w^2 follows from the formulas $\lambda_{0,1}(w^1) = \lambda_{0,1}(w^2) = 0$ and $\lambda_{0,1}(w) \neq 0$.

Set $w = w_{a,b,c}^4$. Formula (12.1.2) with $\sigma(1) = 2, \sigma(2) = 3, \sigma(3) = 1$ yields

$$\lambda(w) = abcb_{\bullet}c_{\bullet}a_{\bullet} + a(1 - bb_{\bullet})c_{\bullet}a_{\bullet} + ab(1 - cc_{\bullet})a_{\bullet} + 1 - aa_{\bullet}.$$

Therefore $\lambda_{0,0}(w) = 1$, and

$$\lambda_{1,0}(w) = ac_{\bullet}a_{\bullet} - abcc_{\bullet}a_{\bullet}, \quad \lambda_{0,1}(w) = aba_{\bullet} - abb_{\bullet}c_{\bullet}a_{\bullet}, \quad \lambda_{1,1}(w) = abcb_{\bullet}c_{\bullet}a_{\bullet} - aa_{\bullet}.$$

If w is contractible, then $\lambda_{1,0}(w) = 0$. This implies $c = \tau(b)$, excluded by the 4-regularity. For a 4-regular triple a, b, c , we can recover a, b, c from $\lambda_{1,0}(w)$. Indeed, $x = z_a$ is the only element of Π such that $(\lambda_{1,0}(w))^{x,y} = 1$ for some $y \in \Pi$. This recovers a . Similarly, $x' = z_a z_b z_c$ is the only element of Π such that $(\lambda_{1,0}(w))^{x',y} = -1$ for some $y \in \Pi$. Knowing x, x' , we recover $x^{-1}x' = z_b z_c$ and the letters b, c . That w is not homotopic to a nanoword of length 4 follows from the formula $\lambda_{0,0}(w) = 1$ and the fact that a nanoword of length 4 with $\lambda_{0,0} = 1$ must be contractible. That w is not homotopic to a nanoword of type w^1 or w^2 follows from the formulas $\lambda_{0,1}(w^1) = \lambda_{0,1}(w^2) = 0$ and $\lambda_{0,1}(w) \neq 0$ (in the 4-regular case). That w is not homotopic to a nanoword of type w^3 follows from the formulas $\lambda_{1,0}(w^3) = 0$ and $\lambda_{1,0}(w) \neq 0$.

Set $w = w_{a,b,c}^5 = ABACBC$. There are 5 paths in the graph Γ_w contributing to $\lambda(w)$. This gives

$$\lambda(w) = (1 - aa_{\bullet})(1 - cc_{\bullet}) + aba_{\bullet}(1 - cc_{\bullet}) + a(1 - bb_{\bullet})c_{\bullet} + (1 - aa_{\bullet})cb_{\bullet}c_{\bullet} + aba_{\bullet}cb_{\bullet}c_{\bullet}.$$

Therefore

$$\lambda_{0,0}(w) = 1 - abb_{\bullet}c_{\bullet} + aa_{\bullet}cc_{\bullet}, \quad \lambda_{1,0}(w) = cb_{\bullet}c_{\bullet} - aba_{\bullet}cc_{\bullet},$$

$$\lambda_{0,1}(w) = aba_{\bullet} - aa_{\bullet}cb_{\bullet}c_{\bullet}, \quad \lambda_{1,1}(w) = ac_{\bullet} - cc_{\bullet} - aa_{\bullet} + aba_{\bullet}cb_{\bullet}c_{\bullet}.$$

Then $\psi(\lambda_{1,0}(w)) = z_c \otimes z_b z_c - z_a z_b z_c \otimes z_a z_c$. Observe that $z_c = z_a z_b z_c \Leftrightarrow a = \tau(b)$ and $z_b z_c = z_a z_c \Leftrightarrow b = a$. Thus if $\lambda_{1,0}(w) = 0$, then $\tau(a) = a = b$. A similar analysis shows that if $\lambda_{0,1}(w) = 0$, then $\tau(b) = b = c$. Therefore if w is contractible, then the triple a, b, c is singular.

Now we show how to recover a, b, c from $w = w_{a,b,c}^5$ provided w is non-contractible. Suppose first that $\psi(\lambda_{1,0}(w)) \neq 0$. Then $x = z_c$ is the only element of Π such that $(\lambda_{1,0}(w))^{x,y} = 1$ for some $y \in \Pi$. This y is unique and equal to $z_b z_c$. This recovers c and b . Then it is easy to recover a from $\psi(\lambda_{1,0}(w))$. If $\psi(\lambda_{1,0}(w)) = 0$, then $\tau(a) = a = b$ and $\psi(\lambda_{1,1}(w)) = a \otimes (c_{\bullet} - a_{\bullet})$. Since w is non-contractible, $a \neq c$. Thus we can recover a and c from $\lambda_{1,1}(w)$.

If w is homotopic to a nanoword w^1 , then $\lambda_{0,1}(w) = \lambda_{1,0}(w) = 0$. As shown above, this implies that the triple a, b, c is singular.

If w is homotopic to a nanoword w^2 , then $\lambda_{0,1}(w) = 0$ and $\lambda_{0,0}(w)$ is a monomial. The latter implies that $abb_{\bullet}c_{\bullet} = 1$ or $abb_{\bullet}c_{\bullet} = aa_{\bullet}cc_{\bullet}$. Applying ψ , we easily deduce that in both cases $a = c$. The equality $\lambda_{0,1}(w) = 0$ implies that $\tau(b) = b = c$. Hence the triple a, b, c is singular.

If w is homotopic to a nanoword w^3 , then $\lambda_{1,0}(w) = 0$ and $\lambda_{0,0}(w)$ is a monomial. The former implies that $a = b = \tau(b)$ and the latter implies that $a = c$. Hence the triple a, b, c is singular.

If w is homotopic to a nanoword w^4 , then $\lambda_{0,0}(w) = 1$. Hence $abb_{\bullet}c_{\bullet} = aa_{\bullet}cc_{\bullet}$. Multiplying on the left by $\tau(a)$ and on the right by $(\tau(c))_{\bullet}$, we obtain $bb_{\bullet} = a_{\bullet}c$. Applying ψ , we easily deduce that $a = b = c$. For $a = b = c$, the nanowords

$w = ABACBC$ and $w_{a,b,c}^4 = ABCBCA$ are related by the homotopy

$$ABACBC \mapsto \underline{A} \underline{D} \underline{E} \underline{B} \underline{A} \underline{C} \underline{B} \underline{E} \underline{D} \underline{C} \mapsto \underline{D} \underline{A} \underline{E} \underline{B} \underline{C} \underline{A} \underline{B} \underline{E} \underline{C} \underline{D} \mapsto \underline{D} \underline{A} \underline{C} \underline{A} \underline{C} \underline{D}$$

where $|D| = a$, $|E| = \tau(a)$ (the nanoword $DACACD$ is isomorphic to $ABCBCA$).

It remains to show that w is not homotopic to a nanoword of length 4. Suppose first that $a \neq c$ or $c \neq \tau(b)$. If w is homotopic to a nanoword of length 4, then $\lambda_{1,1}(w) = 0$ and hence

$$\psi(\lambda_{1,1}(w)) = z_a \otimes z_c - z_c \otimes z_c - z_a \otimes z_a + z_a z_b z_c \otimes z_a z_b z_c = 0.$$

The term $z_a \otimes z_c$ must cancel with either $z_c \otimes z_c$ or $z_a \otimes z_a$. In both cases $a = c$. The term $z_a z_b z_c \otimes z_a z_b z_c$ must then cancel with $z_a \otimes z_a$. This is possible only if $c = \tau(b)$ which contradicts our assumptions.

Consider the remaining case $a = c = \tau(b)$. By the 5-regularity, $a \neq \tau(a)$. We prove here the following weaker claim: if w is homotopic to a nanoword $KLKL$, then $|K| = |L| = a$. That this is also impossible will be proven in Sect. 16.5. To prove the weaker claim, set $k = |K|$, $l = |L|$. Then $\lambda_{1,0}(KLKL) = k - kk_{\bullet} l_{\bullet}$. Since $a = c = \tau(b)$, we have $\lambda_{1,0}(w) = cb_{\bullet} c_{\bullet} - aba_{\bullet} cc_{\bullet} = a - aa_{\bullet} a_{\bullet}$. The equality $k - kk_{\bullet} l_{\bullet} = a - aa_{\bullet} a_{\bullet}$ is possible in only two cases: when $k = l = a$ and when $k = \tau(l)$, $a = \tau(a)$. The latter is excluded by the 5-regularity. Therefore $k = l = a$. $\square$

Corollary 13.2.2. *For $i = 1, \dots, 5$ and any i -regular triple $a, b, c \in \alpha$, we have $\|w_{a,b,c}^i\| = 3$.*

Corollary 13.2.3. *A nanoword of length ≤ 6 is homotopically (skew-) symmetric if and only if it is contractible or (skew-) symmetric.*

One needs to prove only that a homotopically (skew-) symmetric nanoword of length ≤ 6 is contractible or (skew-) symmetric. This follows from the homotopy classification above. Note that $w_{a,b,c}^i$ is symmetric if and only if $i \in \{1, 5\}$, $a = c$ or $i = 4$, $b = c$. The nanoword $w_{a,b,c}^i$ is skew-symmetric if and only if $i \in \{1, 5\}$, $\tau(a) = c$, $\tau(b) = b$ or $i = 4$, $\tau(a) = a$, $\tau(b) = c$.

14. α -QUANDLES AND α -KEIS

We introduce α -quandles and α -keis generalizing the classical quandles and keis. Their connections with words will be discussed in the next section.

14.1. Keis and quandles. Keis were introduced by M. Takasaki in 1942 as abstractions of symmetric transformations. A more general notion of a quandle was introduced in the 1980's by D. Joyce, S. Matveev, and E. Brieskorn independently, see [Ka] for a survey. A *quandle* is a non-empty set X with binary operation $(x, y) \mapsto x * y$ such that $x * x = x$, $(x * y) * z = (x * z) * (y * z)$, for all $x, y, z \in X$, and the mapping $x \mapsto x * z : X \rightarrow X$ is a bijection for any $z \in X$. The identity $(x * y) * z = (x * z) * (y * z)$ means that the bijection $x \mapsto x * z$ preserves $*$. A quandle is a *kei* if this bijection is involutive for all z that is $(x * z) * z = x$ for all $x, z \in X$. For example, $X = \mathbb{Z}/m\mathbb{Z}$ with $x * y = 2y - x$ is a kei for any $m \geq 1$.

14.2. α -quandles. For a set α , we introduce a notion of an α -quandle. Let X be a non-empty set. Suppose that each $a \in \alpha$ gives rise to a bijection $x \mapsto ax : X \rightarrow X$ and to a binary operation $(x, y) \mapsto x *_a y$ on X . These operations form an α -quandle if the following axioms are satisfied:

- (i) $ax *_a x = x$ for all $a \in \alpha, x \in X$;
- (ii) $a(x *_a y) = ax *_a ay$ for all $a \in \alpha, x, y \in X$;
- (iii) $(x *_a y) *_a z = (x *_a az) *_a (y *_a z)$ for all $a \in \alpha, x, y, z \in X$;
- (iv) the mapping $x \mapsto x *_a z : X \rightarrow X$ is a bijection for any $a \in \alpha, z \in X$.

The operations $\{x \mapsto ax : X \rightarrow X\}_{a \in \alpha}$ and $\{(x, y) \mapsto x *_a y\}_{a \in \alpha}$ are called the *quandle operations*. A morphism $X \rightarrow X'$ of α -quandles is a set-theoretic map $X \rightarrow X'$ commuting with these operations.

To give examples of α -quandles, consider the semi-group $\tilde{\Psi}$ with generators $\{a, a_\bullet\}_{a \in \alpha}$ subject to the relations $aa_\bullet = a_\bullet a$ for all $a \in \alpha$. Suppose that $\tilde{\Psi}$ acts on a group X by group automorphisms $x \mapsto ax$ and $x \mapsto a_\bullet x$ where $x \in X$. Then the automorphisms $x \mapsto ax$ together with the binary operations

$$(14.2.1) \quad x *_a y = y(a_\bullet x)(a_\bullet ay)^{-1} \in X$$

for $x, y \in X$ form an α -quandle. We check the axioms. Axiom (i): $ax *_a x = x(a_\bullet ax)(a_\bullet ax)^{-1} = x$. Axiom (ii):

$$a(x *_a y) = a(y(a_\bullet x)(a_\bullet ay)^{-1}) = (ay)(aa_\bullet x)(aa_\bullet ay)^{-1} = ax *_a ay.$$

Axiom (iii):

$$\begin{aligned} (x *_a az) *_a (y *_a z) &= (az)(a_\bullet x)(a_\bullet aaz)^{-1} *_a z(a_\bullet y)(a_\bullet az)^{-1} \\ &= z(a_\bullet y)(a_\bullet az)^{-1}(a_\bullet az)(a_\bullet a_\bullet x)(a_\bullet a_\bullet aaz)^{-1}((a_\bullet az)(a_\bullet aa_\bullet y)(a_\bullet aa_\bullet az)^{-1})^{-1} \\ &= z(a_\bullet y)(a_\bullet az)^{-1}(a_\bullet az)(a_\bullet a_\bullet x)(a_\bullet a_\bullet aaz)^{-1}(a_\bullet a_\bullet aaz)(a_\bullet a_\bullet ay)^{-1}(a_\bullet az)^{-1} \\ &= z(a_\bullet y)(a_\bullet a_\bullet x)(a_\bullet a_\bullet ay)^{-1}(a_\bullet az)^{-1} = z a_\bullet (x *_a y)(a_\bullet az)^{-1} = (x *_a y) *_a z. \end{aligned}$$

Axiom (iv) follows from the assumption that $x \mapsto a_\bullet x : X \rightarrow X$ is a bijection.

In particular, any left module over the semi-group ring $\mathbb{Z}\tilde{\Psi}$ is an α -quandle with quandle operations $x \mapsto ax$ and $x *_a y = a_\bullet x + (1 - a_\bullet a)y$.

Note that if $\alpha = \{a\}$ is a 1-element set and $ax = x$ for all x , then an α -quandle is simply a quandle.

14.3. α -keis. Consider a set α with involution τ . An α -kei is an α -quandle X such that

- (v) $a\tau(a)x = x$ for all $x \in X, a \in \alpha$ and
- (vi) $(x *_a y) *_\tau(a) ay = x$ for all $x, y \in X, a \in \alpha$.

These two axioms strengthen Axiom (iv) above: they imply that for any $y \in X$, the mappings $x \mapsto x *_a y : X \rightarrow X$ and $x \mapsto x *_\tau(a) ay : X \rightarrow X$ are inverse to each other.

The quandle operations $\{x \mapsto ax : X \rightarrow X\}_{a \in \alpha}$ and $\{(x, y) \mapsto x *_a y\}_{a \in \alpha}$ in a kei X are called the *kei operations*. Morphisms of α -keis are their morphisms as α -quandles. Isomorphism of α -keis is denoted by $\approx$.

Recall the ring $\Lambda = \mathbb{Z}\Psi$ defined in Sect. 10.3. Any left Λ -module X becomes an α -kei with kei operations $x \mapsto ax$ and $x *_a y = a_\bullet x + (1 - a_\bullet a)y$ for $x, y \in X$.

Axioms (i) – (iv) were checked above, Axiom (v) follows from the definition of Λ and Axiom (vi) follows from the formulas

$$\begin{aligned} (x *_a y) *_{\tau(a)} ay &= \tau(a) \bullet (x *_a y) + (1 - \tau(a) \bullet \tau(a)) ay \\ &= \tau(a) \bullet a \bullet x + \tau(a) \bullet y - \tau(a) \bullet a \bullet ay + ay - \tau(a) \bullet \tau(a) ay = x. \end{aligned}$$

The α -keis obtained by this construction from Λ -modules are said to be *abelian*.

When α is a 1-element set and $ax = x$ for all x , the notion of an α -kei is equivalent to the one of a kei.

14.4. Presentations by generators and relations. For an α -kei X , a set $S \subset X$ *generates* X if all elements of X can be obtained from elements of S using the kei operations. Any set S generates a unique *free α -kei* $X(S)$ characterized by the condition that every set-theoretic map from S to an α -kei X extends to a morphism of α -keis $X(S) \rightarrow X$. The elements of $X(S)$ are obtained from elements of S using the kei operations modulo the identities imposed by Axioms (i) – (iii), (v), (vi).

As in the theory of groups, we can present an α -kei by a set of generators S and a set of relations R . A *relation* is a pair p, q of elements of $X(S)$ which we write as an equality $p = q$. The α -kei presented by S and R is obtained by quotienting $X(S)$ by all the relations from R and all their corollaries. For example, pick $a, b \in \alpha$ and consider the α -kei X with generators x, y subject to one defining relation $x *_b b\tau(a)y = (bx *_a y) *_{\tau(b)} aax$. This means that every element of X can be obtained from x, y using the kei operations, that the relation in question is satisfied, and that all other relations between x and y are corollaries of this one and the axioms of an α -kei. Following a similar train of ideas, one can define free products of α -keis. We leave the details to the reader, cf. [FR] for presentations of quandles.

14.5. Abelianization. Each α -kei X gives rise to a Λ -module with generators $\{[x]\}_{x \in X}$ subject to the “commutation relations” $[ax] = a[x]$ and $[x *_a y] = a \bullet [x] + (1 - a \bullet a)[y]$ for all $x, y \in X, a \in \alpha$. This module is called the *abelianization* of X and denoted X^{ab} . It can be described by the following universal property: any kei morphism from X to an abelian α -kei Y factors uniquely as a composition of the mapping $x \mapsto [x] : X \rightarrow X^{ab}$ and a Λ -homomorphism $X^{ab} \rightarrow Y$.

Given a presentation of X by generators S and relations R we can compute X^{ab} in terms of generators and relations. This module is generated by the set $\{[s]\}_{s \in S}$ and each relation $p = q$ from R gives rise to a relation obtained by expanding both p and q as linear combinations of the vectors $\{[s]\}_{s \in S}$ via the commutation relations. In particular, for the free α -kei $X(S)$ generated by S , we have $X(S)^{ab} = \oplus_{s \in S} \Lambda[s]$. For each $v \in X(S)$, we have a unique expansion $[v] = \sum_{s \in S} \lambda_s(v)[s]$ with $\lambda_s(v) \in \Lambda$.

14.6. Marked α -keis. An α -kei X endowed with an ordered pair of distinguished elements $v_-, v_+ \in X$ is *marked*. The elements v_-, v_+ are called the *input* and the *output*, respectively. By *(iso)morphisms* of marked α -keis, we mean kei

(iso)morphisms preserving the input and the output. The abelianization of a marked α -kei (X, v_-, v_+) is the marked Λ -module $(X^{ab}, [v_-] \in X^{ab}, [v_+] \in X^{ab})$.

Given a marked α -kei X we define a marked α -kei $\overline{X}$ to be the same set X with the same input and output and new kei operations $ax := \tau(a)x$, $x *_a y := x *_{\tau(a)} y$ for $x, y \in X, a \in \alpha$. Clearly, $\overline{\overline{X}} = X$. A presentation of X by generators S and relations R yields a presentation of $\overline{X}$ by generators S and relations $\overline{R}$ where the relations in $\overline{R}$ are obtained from those in R by replacing each letter $a \in \alpha$ appearing in these relations by $\tau(a)$. We define a marked α -kei X^- to be $\overline{X}$ with permuted input and output. Clearly, $(X^-)^- = X$.

We point out a simple numerical invariant of a finitely generated marked α -kei $\mathcal{K}$. Fix a marked α -kei X that is finite as a set. Then the set of marked morphisms $\mathcal{K} \rightarrow X$ is finite. The number of such morphisms is an isomorphism invariant of $\mathcal{K}$.

14.7. Reconstruction. Given an α -quandle X and a set $\alpha_0 \subset \alpha$, we can restrict the quandle operations in X to only those a which belong to α_0 . This gives an α_0 -quandle Y coinciding with X as a set and called the *restriction* of X to α_0 . If X is an α -kei, then for any $a \in \alpha_0 \cap \tau(\alpha_0)$ and any $x, y \in Y$, we have $a\tau(a)x = x$ and $(x *_a y) *_{\tau(a)} ay = x$. An α_0 -quandle Y satisfying the latter conditions is said to be *compatible* with τ .

Lemma 14.7.1. *Let α_0 be a subset of α such that $\alpha_0 \cup \tau(\alpha_0) = \alpha$. Let Y be an α_0 -quandle compatible with τ . Then there is a unique α -kei X whose restriction to α_0 is Y .*

Proof. Clearly, $X = Y$ as a set and an α_0 -quandle. We need only to define the operations $x \mapsto ax$ and $(x, y) \mapsto x *_a y$ for $a \in \alpha - \alpha_0$. By assumption, $\tau(a) \in \alpha_0$. By the definition of an α_0 -quandle, the mappings $x \mapsto \tau(a)x : X \rightarrow X$ and $x \mapsto x *_{\tau(a)} ay : X \rightarrow X$ are bijective. The axioms of an α -kei show that we must take their inverses as the mappings $x \mapsto ax$ and $x \mapsto x *_a y$, respectively. Thus $z = ax$ is the only element of X such that $\tau(a)z = x$ and $t = x *_a y$ is the only element of X such that $t *_{\tau(a)} ay = x$. We show that this makes X into an α -kei.

We first check the axioms of an α -quandle. It suffices to check them for $a \in \alpha - \alpha_0$. Axiom (i): to verify that $ax *_a x = x$ it suffices to check that $x *_{\tau(a)} ax = ax$. This holds since

$$x *_{\tau(a)} ax = a\tau(a)(x *_{\tau(a)} ax) = a(\tau(a)x *_{\tau(a)} \tau(a)ax) = a(\tau(a)x *_{\tau(a)} x) = ax.$$

To check (ii), set $t = x *_a y$. The required equality $at = ax *_a ay$ is equivalent to $at *_{\tau(a)} aay = ax$. We have

$$at *_{\tau(a)} aay = a\tau(a)(at *_{\tau(a)} aay) = a(\tau(a)at *_{\tau(a)} \tau(a)aay) = a(t *_{\tau(a)} ay) = ax$$

by the choice of t . To check (iii), set $t = (x *_a y) *_a z$. The required equality $t = (x *_a az) *_a (y *_a z)$ is equivalent to $t *_{\tau(a)} a(y *_a z) = x *_a az$. The latter is equivalent to

$$(t *_{\tau(a)} a(y *_a z)) *_{\tau(a)} aaz = x.$$

By Axiom (iii) applied to $\tau(a) \in \alpha_0$,

$$\begin{aligned} (t *_{\tau(a)} a(y *_{\alpha} z)) *_{\tau(a)} aaz &= (t *_{\tau(a)} \tau(a)aaz) *_{\tau(a)} (a(y *_{\alpha} z) *_{\tau(a)} aaz) \\ &= (t *_{\tau(a)} az) *_{\tau(a)} ((ay *_{\alpha} az) *_{\tau(a)} aaz) = (x *_{\alpha} y) *_{\tau(a)} ay = x. \end{aligned}$$

Axiom (iv) follows directly from the definitions.

To see that X is an α -kei, we must check that $(x *_{\alpha} y) *_{\tau(a)} ay = x$ for all $x, y \in X, a \in \alpha$. This follows from the definition of $*_{\alpha}$ for $a \in \alpha - \alpha_0$ and from the compatibility assumption for $a \in \alpha_0 \cap \tau(\alpha_0)$. It remains to consider the case where $a \in \alpha_0$ and $\tau(a) \notin \alpha_0$. By the definition of $*_{\tau(a)}$, we know that $t = (x *_{\alpha} y) *_{\tau(a)} ay$ is the only element of X such that $t *_{\alpha} \tau(a)ay = x *_{\alpha} y$. Clearly, $t = x$ is such an element. Therefore $(x *_{\alpha} y) *_{\tau(a)} ay = x$. $\square$

This lemma establishes a bijective correspondence between (the isomorphisms classes of) α -keis and α_0 -quandles compatible with τ . If $\alpha_0 \cap \tau(\alpha_0) = \emptyset$, then the compatibility condition is empty and we obtain a bijective correspondence between α -keis and α_0 -quandles.

15. KEIS OF NANOWORDS

With each nanoword w over α and a τ -invariant set $\beta \subset \alpha$ we associate a marked α -kei $\mathcal{K}_{\beta}(w)$. It is preserved under homotopy moves on w .

15.1. The α -kei $\mathcal{K}_{\beta}(w)$. Consider a nanoword $(\mathcal{A}, w : \hat{n} \rightarrow \mathcal{A})$ over α . As in Sect. 8.1, for a letter $A \in \mathcal{A}$, we denote by i_A (resp. j_A) the minimal (resp. the maximal) element of the 2-element set $\omega^{-1}(A) \subset \hat{n}$. The α -kei $\mathcal{K}_{\beta}(w)$ is generated by $n + 1$ symbols $x_0, x_1, \dots, x_n$ subject to the following n defining relations: for any $A \in \mathcal{A}$ such that $a = |A| \in \beta$,

$$x_{i_A} = a x_{i_A-1}, \quad x_{j_A} = x_{j_A-1} *_{\alpha} x_{i_A-1},$$

and for any $A \in \mathcal{A}$ such that $a = |A| \in \alpha - \beta$,

$$x_{i_A} = x_{i_A-1} *_{\alpha} x_{j_A-1}, \quad x_{j_A} = a x_{j_A-1}.$$

We take x_0 as the input and x_n as the output in $\mathcal{K}_{\beta}(w)$. Comparing this definition with the one of the marked module $K_{\beta}(w)$ we obtain that

$$K_{\beta}(w) = (\mathcal{K}_{\beta}(w))^{ab}.$$

It is clear that if $w = w_1 w_2$ is a product of two nanowords w_1, w_2 , then $\mathcal{K}(w)$ is the free product of $\mathcal{K}(w_1)$ and $\mathcal{K}(w_2)$ quotiented by the relation (the output of $\mathcal{K}(w_1)$) = (the input of $\mathcal{K}(w_2)$).

It follows from the definitions that $\mathcal{K}_{\beta}(\overline{w}) \approx \overline{\mathcal{K}_{\beta}(w)}$. The isomorphism is the identity $x_i \mapsto x_i$ on the generators. Similarly, $\mathcal{K}_{\beta}(w^-) \approx (\mathcal{K}_{\alpha-\beta}(w))^-$. The latter isomorphism is induced by the permutation of the generators $x_i \mapsto x_{n-i}$. The main property of $\mathcal{K}_{\beta}(w)$ is contained in the next lemma.

Lemma 15.1.1. *The isomorphism type of the marked α -kei $\mathcal{K}_{\beta}(w)$ is a homotopy invariant of w .*

Proof. If w is replaced by an isomorphic nanoword, then $\mathcal{K}_\beta(w)$ is replaced by an isomorphic marked α -kei. Consider the first homotopy move $w = xAAy \mapsto xy = v$. Set $a = |A| \in \alpha$ and let $i = i_A, i+1 = j_A$ be the indices numerating the entries of A in w . If $a \in \beta$, then the generator x_i of $\mathcal{K}_\beta(w)$ is involved in two relations $x_i = ax_{i-1}$ and $x_{i+1} = x_i *_a x_{i-1}$. We can exclude x_i from the set of generators using the first relation and replace the second relation with $x_{i+1} = ax_{i-1} *_a x_{i-1}$. By Axiom (i), this may be rewritten as $x_{i+1} = x_{i-1}$. Thus we can further remove x_{i+1} from the set of generators and replace it in the remaining relations by x_{i-1} . The resulting set of generators and relations is a presentation of $\mathcal{K}_\beta(v)$.

If $a \in \alpha - \beta$, then

$$(15.1.1) \quad \mathcal{K}_\beta(w) \approx (\mathcal{K}_{\alpha-\beta}(w^-))^- \approx (\mathcal{K}_{\alpha-\beta}(v^-))^- \approx \mathcal{K}_\beta(v)$$

where the second isomorphism follows from the inclusion $a \in \alpha - \beta$ and the result of the previous paragraph. For completeness, we give a direct proof of the isomorphism $\mathcal{K}_\beta(w) \approx \mathcal{K}_\beta(v)$. The generator x_i of $\mathcal{K}_\beta(w)$ is involved in two relations $x_{i+1} = ax_i$ and $x_i = x_{i-1} *_a x_i$. The first formula is equivalent to $x_i = \tau(a)x_{i+1}$. We can exclude x_i via this relation and rewrite the second relation as $\tau(a)x_{i+1} = x_{i-1} *_a \tau(a)x_{i+1}$. Note that for any elements x, y of an α -kei,

$$\tau(a)x = y *_a \tau(a)x \iff x = a(y *_a \tau(a)x) \iff x = ay *_a x \iff x = y.$$

Therefore we can rewrite the relation $\tau(a)x_{i+1} = x_{i-1} *_a \tau(a)x_{i+1}$ as $x_{i+1} = x_{i-1}$. Thus we can further remove x_{i+1} from the set of generators and replace it in the remaining relations by x_{i-1} . The resulting set of generators and relations is a presentation of $\mathcal{K}_\beta(v)$.

Consider the second homotopy move $w = xAB yBAz \mapsto xyz = v$. Set $a = |A| \in \alpha$ and let $i = i_A, j = j_A \geq i+3$ be the indices numerating the entries of A in w . By assumption, $|B| = \tau(a)$. If $a \in \beta$, then $\tau(a) \in \beta$ and the letters A, B give rise to four relations $x_i = ax_{i-1}, x_{i+1} = \tau(a)x_i$ and $x_{j-1} = x_{j-2} *_a x_i, x_j = x_{j-1} *_a x_{i-1}$. The first two relations imply that $x_{i+1} = x_{i-1}$. We exclude x_i, x_{i+1} from the set of generators replacing them in all other relations with ax_{i-1}, x_{i-1} respectively. The resulting relation $x_{j-1} = x_{j-2} *_a ax_{i-1}$ allows us to exclude x_{j-1} from the set of generators and to rewrite the relation $x_j = x_{j-1} *_a x_{i-1}$ as $x_j = (x_{j-2} *_a ax_{i-1}) *_a x_{i-1}$. The latter is equivalent to $x_j = x_{j-2}$. Thus we can further remove x_j from the set of generators and replace it in the remaining relations by x_{j-2} . The resulting set of generators and relations is a presentation of $\mathcal{K}_\beta(v)$. The case $a \in \alpha - \beta$ follows from the case $a \in \beta$ by (15.1.1).

Consider the third homotopy move $w = xAB yACzBCt \mapsto xBA yCAzCBt = v$. Set $a = |A| = |B| = |C| \in \alpha$. Let $i = i_A, j = j_A, k = j_B$ be the indices numerating the entries of A in w and the second entry of B in w . If $a \in \beta$, then the letters A, B, C give rise to six relations in $\mathcal{K}_\beta(w)$, namely, $x_i = ax_{i-1}$ and

$$x_{i+1} = ax_i, x_j = x_{j-1} *_a x_{i-1}, x_{j+1} = ax_j, x_k = x_{k-1} *_a x_i, x_{k+1} = x_k *_a x_j.$$

We exclude x_i, x_j, x_k using the first, third, and fifth relations rewriting the other 3 relations as

$$(15.1.2) \quad x_{i+1} = aa x_{i-1}, x_{j+1} = a(x_{j-1} *_a x_{i-1}),$$

$$x_{k+1} = (x_{k-1} *_a ax_{i-1}) *_a (x_{j-1} *_a x_{i-1}).$$

Similarly, A, B, C give rise to 6 relations in $\mathcal{K}_\beta(v)$, namely, $x_i = ax_{i-1}$ and

$$x_{i+1} = ax_i, x_j = ax_{j-1}, x_{j+1} = x_j *_a x_i, x_k = x_{k-1} *_a x_{j-1}, x_{k+1} = x_k *_a x_{i-1}.$$

We exclude x_i, x_j, x_k using the first, third, and fifth relations rewriting the other 3 relations as

$$x_{i+1} = aax_{i-1}, x_{j+1} = ax_{j-1} *_a ax_{i-1}, x_{k+1} = (x_{k-1} *_a x_{j-1}) *_a x_{i-1}.$$

These relations are equivalent to the relations (15.1.2) modulo the axioms of an α -kei. The case $a \in \alpha - \beta$ follows from the case $a \in \beta$ by (15.1.1). $\square$

15.2. The invariant $v_+(w)$. For $\beta = \alpha$, the defining relations of the α -kei $\mathcal{K}_\beta(w) = \mathcal{K}_\alpha(w)$ consecutively express the generators via the previous ones. Therefore $\mathcal{K}_\alpha(w)$ is a free α -kei generated by the input v_- . We identify $\mathcal{K}_\alpha(w)$ with the free α -kei $X(s)$ with one generator s via $v_- = s$. The output $v_+(w) = v_+ \in \mathcal{K}_\alpha(w) = X(s)$ is a homotopy invariant of w . It includes the polynomials $\lambda'(w), \lambda(w) \in \Lambda$ studied in Sect. 12.1 via $[v_+(w)] = \lambda'(w)[s] \in (X(s))^{ab} = \Lambda[s]$.

Since $X(s)$ is a free α -kei, the element $v_+(w) \in X(s)$ determines a unique kei morphism $\phi(w) : X(s) \rightarrow X(s)$ such that $\phi(w)(s) = v_+(w)$. This morphism is a homotopy invariant of w . If $w = w_1 w_2$ is a product of two nanowords, then $\phi(w) = \phi(w_1) \phi(w_2)$. Indeed, both sides are kei endomorphisms of $X(s)$ transforming s into $v_+(w)$.

We finish this section by showing how to compute $v_+(\overline{w})$ from $v_+(w)$. Given two α -keis X, X' , we call a mapping $f : X \rightarrow X'$ a *kei anti-morphism* if $f(ax) = \tau(a)x$ and $f(x *_a y) = f(x) *_a f(y)$ for all $x, y \in X, a \in \alpha$. In other words, a kei anti-morphism $X \rightarrow X'$ is a kei morphism $X \rightarrow \overline{X'}$. It is obvious that $\overline{X(s)}$ is a free α -kei generated by s and therefore there is a unique kei anti-morphism $\iota : X(s) \rightarrow X(s)$ preserving s . Clearly, ι is an involution. It follows from the definitions that $v_+(\overline{w}) = \iota(v_+(w))$.

16. CASE OF FREE τ

Throughout this section we suppose that τ is fixed-point-free and fix a set $\alpha_0 \subset \alpha$ meeting each orbit of τ in one element. We derive from α -keis of nanowords a simpler invariant, called the characteristic sequence. It is used to accomplish the homotopy classification of nanowords of length ≤ 6 .

16.1. The α -kei F . We first construct an α -kei F needed for the sequel. Recall from Sect. 10.3 the group $\Psi = \Psi_\alpha$ with generators $\{a, a_\bullet\}_{a \in \alpha}$ and defining relations $aa_\bullet = a_\bullet a, a\tau(a) = a_\bullet \tau(a)_\bullet = 1$ for $a \in \alpha$. Let F be the free group (of countable rank) freely generated by the set Ψ . The generator of F corresponding to $\psi \in \Psi$ is denoted $\underline{\psi}$. Note that $\underline{\psi}$ and $\underline{\psi}^{-1}$ are two independent generators of F for $\psi \neq 1$ and $\underline{1}$ is a non-trivial generator of F where 1 is the unit of Ψ . A typical element of F has the form $(\underline{\psi_1})^{\varepsilon_1} \cdots (\underline{\psi_m})^{\varepsilon_m}$ where $m \geq 0, \psi_1, \dots, \psi_m \in \Psi$, and $\varepsilon_1, \dots, \varepsilon_m \in \{\pm 1\}$.

The left action of Ψ on itself extends to a group action of Ψ on F . In particular, the generators $a, a_\bullet \in \Psi$ act on F by the group automorphisms

$$\begin{aligned} a((\underline{\psi_1})^{\varepsilon_1} \dots (\underline{\psi_m})^{\varepsilon_m}) &= (\underline{a\psi_1})^{\varepsilon_1} \dots (\underline{a\psi_m})^{\varepsilon_m}, \\ a_\bullet((\underline{\psi_1})^{\varepsilon_1} \dots (\underline{\psi_m})^{\varepsilon_m}) &= (\underline{a_\bullet\psi_1})^{\varepsilon_1} \dots (\underline{a_\bullet\psi_m})^{\varepsilon_m}. \end{aligned}$$

By Sect. 14.3, this induces a structure of an α_0 -quandle on F with quandle operations $x \mapsto ax$ and $x *_a y = y(a_\bullet x)(a_\bullet ay)^{-1}$ for $a \in \alpha_0$. By Lemma 14.7.1, this extends uniquely to an α -kei structure on F . The compatibility condition in Lemma 14.7.1 is empty since $\alpha_0 \cap \tau(\alpha_0) = \emptyset$. Note that the resulting α -kei structure on F , generally speaking, depends on the choice of α_0 . The next lemma gives explicit formulas for the kei operations in F determined by elements of $\alpha - \alpha_0$.

Lemma 16.1.1. *For $a \in \alpha_0$, $\psi_1, \dots, \psi_m \in \Psi$, and $\varepsilon_1, \dots, \varepsilon_m \in \{\pm 1\}$,*

$$\tau(a)((\underline{\psi_1})^{\varepsilon_1} \dots (\underline{\psi_m})^{\varepsilon_m}) = (\underline{a^{-1}\psi_1})^{\varepsilon_1} \dots (\underline{a^{-1}\psi_m})^{\varepsilon_m} = (\underline{\tau(a)\psi_1})^{\varepsilon_1} \dots (\underline{\tau(a)\psi_m})^{\varepsilon_m}.$$

For $a \in \alpha_0$, $x, y \in F$,

$$(16.1.1) \quad x *_{\tau(a)} y = (a_\bullet^{-1} a^{-1} y)^{-1} (a_\bullet^{-1} x) y \in F.$$

Proof. Recall from the proof of Lemma 14.7.1 that $z = \tau(a)x \in F$ and $t = x *_{\tau(a)} y$ are uniquely determined from the equations $az = x$ and $t *_a \tau(a)y = x$, respectively. Observe that

$$a((\underline{\tau(a)\psi_1})^{\varepsilon_1} \dots (\underline{\tau(a)\psi_m})^{\varepsilon_m}) = (\underline{a\tau(a)\psi_1})^{\varepsilon_1} \dots (\underline{a\tau(a)\psi_m})^{\varepsilon_m} = (\underline{\psi_1})^{\varepsilon_1} \dots (\underline{\psi_m})^{\varepsilon_m}.$$

This implies the first claim of the lemma. We have

$$\begin{aligned} (a_\bullet^{-1} a^{-1} y)^{-1} (a_\bullet^{-1} x) y *_a \tau(a)y &= \tau(a)y a_\bullet((a_\bullet^{-1} a^{-1} y)^{-1} (a_\bullet^{-1} x) y) (a_\bullet a\tau(a)y)^{-1} \\ &= \tau(a)y (a_\bullet a_\bullet^{-1} a^{-1} y)^{-1} (a_\bullet a_\bullet^{-1} x) a_\bullet y (a_\bullet y)^{-1} = \tau(a)y (a^{-1} y)^{-1} x = x \end{aligned}$$

where we use that $a\tau(a)y = y$ and $a^{-1}y = \tau(a)y$. This implies (16.1.1). $\square$

16.2. Characteristic sequences. We can use the α -kei F constructed above to study the free α -kei $X(S)$ generated by a set S . By the definition of $X(S)$, there is a unique kei morphism $f : X(S) \rightarrow F$ sending S to $\underline{1} \in F$. For $v \in X(S)$, we have $f(v) = (\underline{\psi_1})^{\varepsilon_1} \dots (\underline{\psi_m})^{\varepsilon_m}$ for some $\varepsilon_1, \dots, \varepsilon_m \in \{\pm 1\}$ and $\psi_1, \dots, \psi_m \in \Psi$. The sequence $(\varepsilon_1\psi_1, \dots, \varepsilon_m\psi_m)$ is called a *characteristic sequence* of v . It is well defined up to insertion/deletion of pairs of consecutive terms $+\psi, -\psi$ or $-\psi, +\psi$ with $\psi \in \Psi$. Deleting all such pairs we obtain a unique *reduced characteristic sequence* of v . It can be efficiently computed. For example, let us compute the reduced characteristic sequence of $v = (bas *_a s) *_b as$ where $a, b \in \alpha, s \in S$. By definition $f(s) = \underline{1}$,

$$\begin{aligned} f(bas) &= ba\underline{1} = \underline{ba}, \quad f(bas *_a s) = f(bas) *_a f(s) = \underline{ba} *_a \underline{1} = \underline{1 a_\bullet ba (a_\bullet a)^{-1}}, \\ f((bas *_a s) *_b as) &= f(bas *_a s) *_b f(as) \\ &= (\underline{1 a_\bullet ba (a_\bullet a)^{-1}}) *_b \underline{a} = \underline{a} \underline{b_\bullet} \underline{b_\bullet a_\bullet ba (b_\bullet a_\bullet a)^{-1} (b_\bullet ba)^{-1}}. \end{aligned}$$

Hence $(a, b_\bullet, b_\bullet a_\bullet ba, -b_\bullet a_\bullet a, -b_\bullet ba)$ is a characteristic sequence of v . It is reduced since $b \neq 1 \in \Psi$.

The following lemma relates the characteristic sequences to the polynomials $\lambda_s(v) \in \Lambda = \mathbb{Z}\Psi$ defined in Sect. 14.5.

Lemma 16.2.1. *Let $(\varepsilon_1\psi_1, \dots, \varepsilon_m\psi_m)$ be a characteristic sequence of $v \in X(S)$. Then*

$$\varepsilon_1\psi_1 + \dots + \varepsilon_m\psi_m = \sum_{s \in S} \lambda_s(v).$$

Proof. The canonical inclusion $\Psi \hookrightarrow \mathbb{Z}\Psi = \Lambda$ extends uniquely to a group homomorphism i from F to the underlying additive group of Λ . In particular,

$$i(f(v)) = i((\underline{\psi_1})^{\varepsilon_1} \cdots (\underline{\psi_m})^{\varepsilon_m}) = \varepsilon_1\psi_1 + \dots + \varepsilon_m\psi_m.$$

The ring Λ being a left Λ -module acquires the structure of an abelian α -kei. Comparing the definitions of the kei operations in F and Λ we observe that i is a kei morphism. The mapping $if : X(S) \rightarrow \Lambda$ splits therefore as a composition of the projection $x \mapsto [x] : X(S) \rightarrow X(S)^{ab}$ and a Λ -homomorphism $f' : X(S)^{ab} \rightarrow \Lambda$. For any $s \in S$, we have $f'([s]) = if(s) = i(\underline{1}) = 1$. Therefore

$$\varepsilon_1\psi_1 + \dots + \varepsilon_m\psi_m = i(f(v)) = f'([v]) = f'(\sum_{s \in S} \lambda_s(v)[s]) = \sum_{s \in S} \lambda_s(v).$$

□

Lemma 16.2.2. *Let $\tau_{\#} : \Psi \rightarrow \Psi$ be the involutive group automorphism sending the generators $a, a_{\bullet}$ to $\tau(a) = a^{-1}, \tau(a)_{\bullet} = a_{\bullet}^{-1}$, respectively, for $a \in \alpha$. If $(\varepsilon_1\psi_1, \dots, \varepsilon_m\psi_m)$ is a characteristic sequence of $v \in X(S)$, then $(\varepsilon_m\tau_{\#}(\psi_m), \dots, \varepsilon_1\tau_{\#}(\psi_1))$ is a characteristic sequence of $\iota(v)$ where ι is the unique kei anti-automorphism of $X(S)$ preserving S element-wise.*

Proof. Denote by $\tilde{\tau}$ the unique group anti-automorphism of F extending $\tau_{\#} : \Psi \rightarrow \Psi$. Observe that $\tilde{\tau}(ax) = a^{-1}\tilde{\tau}(x)$ and $\tilde{\tau}(a_{\bullet}x) = a_{\bullet}^{-1}\tilde{\tau}(x)$ for $a \in \alpha, x \in F$. Moreover, $\tilde{\tau}$ is a kei anti-morphism. Indeed, for $x, y \in F, a \in \alpha_0$,

$$\begin{aligned} \tilde{\tau}(x *_a y) &= \tilde{\tau}(y(a_{\bullet}x)(a_{\bullet}ay)^{-1}) = (\tilde{\tau}(a_{\bullet}ay))^{-1} \tilde{\tau}(a_{\bullet}x) \tilde{\tau}(y) \\ &= (a_{\bullet}^{-1}a^{-1}\tilde{\tau}(y))^{-1} a_{\bullet}^{-1}\tilde{\tau}(x) \tilde{\tau}(y) = \tilde{\tau}(x) *_{\tau(a)} \tilde{\tau}(y). \end{aligned}$$

The case $a \in \alpha - \alpha_0$ is similar. The compositions $f\iota : X(S) \rightarrow F$ and $\tilde{\tau}f : X(S) \rightarrow F$ are kei anti-morphisms sending S to $\underline{1}$. Therefore $f\iota = \tilde{\tau}f$. The equality $f\iota(v) = \tilde{\tau}f(v)$ implies the claim of the lemma. □

16.3. Characteristic sequences of nanowords. By a *characteristic sequence* of a nanoword w , we mean a characteristic sequence of the element $v_+(w) \in X(s)$ defined in Sect. 15.2. This sequence, say $(\varepsilon_1\psi_1, \dots, \varepsilon_m\psi_m)$, is well defined up to insertion of consecutive terms $+\psi, -\psi$ or $-\psi, +\psi$ with $\psi \in \Psi$. Deleting all such terms we obtain the *reduced characteristic sequence* of w . Lemma 16.2.1 implies that $\lambda'(w) = \varepsilon_1\psi_1 + \dots + \varepsilon_m\psi_m$. Lemma 16.2.2 implies that $(\varepsilon_m\tau_{\#}(\psi_m), \dots, \varepsilon_1\tau_{\#}(\psi_1))$ is a characteristic sequence of $\overline{w}$. By Lemma 15.1.1, the reduced characteristic sequence of a nanoword is a homotopy invariant. Note that the α -kei structure on F and hence the characteristic sequences of nanowords depend on the choice of α_0 .

16.4. Examples. 1. Let $w = ABAB$ with $|A| = a \in \alpha_0, |B| = b \in \alpha_0$. The α -kei $\mathcal{K}_\alpha(w)$ is generated by $v_- = x_0, x_1, \dots, x_4 = v_+$ subject to the relations $x_1 = ax_0, x_2 = bx_1, x_3 = x_2 *_a x_0, x_4 = x_3 *_b x_1$. Thus $v_+ = (bav_- *_a v_-) *_b av_-$ and $v_+(w) = (bas *_a s) *_b as \in X(s)$. By the computations above, the reduced characteristic sequence of w is $(a, b_\bullet, b_\bullet a_\bullet ba, -b_\bullet a_\bullet a, -b_\bullet ba)$.

2. Pick four letters $a, b, c, d \in \alpha$ (possibly coinciding) and consider the nanoword $w = ABCDCDAB$ with $|A| = a, |B| = b, |C| = c, |D| = d$. An inspection shows that if $a \neq \tau(b)$ and $c \neq \tau(d)$, then the α -pairing of w is primitive. Then $\|w\| = 4$ and w is non-contractible. However, for any $a, b \in \alpha_0$ and $c = \tau(b), d = \tau(a)$ we have $f(v_+(w)) = \underline{1}$ (a direct computation). In this case, the characteristic sequence does not distinguish w from a contractible nanoword. This shows that the characteristic sequence is not a faithful invariant of nanowords and does not allow to recover the associated α -pairing.

16.5. End of the proof of Theorem 13.2.1. We must show that the nanowords $ABACBC$ and $ACAC$ with $|A| = |C| = a, |B| = \tau(a) \neq a$ are not homotopic. By Lemma 3.3.1, it suffices to consider the case where the alphabet α consists of only two letters a and $\tau(a) \neq a$. Then τ is fixed-point-free and we can consider the characteristic sequences of nanowords determined by $\alpha_0 = \{a\} \subset \alpha$. The computation above shows that the reduced characteristic sequence of $ACAC$ is $(a, a_\bullet, a^2 a_\bullet^2, -aa_\bullet^2, -a^2 a_\bullet)$. It follows from the definitions that

$$v_+(ABACBC) = (a(s *_a s) *_{\tau(a)} as) *_a (s *_a s).$$

The value of $f : X(s) \rightarrow F$ on $v_+(ABACBC)$ can be easily computed:

$$\begin{aligned} f(s *_a s) &= \underline{1} *_a \underline{1} = \underline{1} \underline{a_\bullet} (\underline{a_\bullet a})^{-1}, \quad f(a(s *_a s)) = \underline{a} \underline{aa_\bullet} (\underline{aa_\bullet a})^{-1}, \\ f(a(s *_a s) *_{\tau(a)} as) &= \underline{a} \underline{aa_\bullet} (\underline{aa_\bullet a})^{-1} *_{\tau(a)} \underline{a} = \left(\underline{a_\bullet^{-1}} \right)^{-1} \underline{a_\bullet^{-1} a} \underline{a} (\underline{aa})^{-1} \underline{a}, \\ f(v_+(ABACBC)) &= \left(\underline{a_\bullet^{-1}} \right)^{-1} \underline{a_\bullet^{-1} a} \underline{a} (\underline{aa})^{-1} \underline{a} *_a \underline{1} \underline{a_\bullet} (\underline{a_\bullet a})^{-1} \\ &= \underline{1} \underline{a_\bullet} (\underline{aa_\bullet})^{-1} (\underline{1})^{-1} \underline{a} \underline{aa_\bullet} (\underline{a^2 a_\bullet})^{-1} \underline{aa_\bullet} \underline{a^2 a_\bullet^2} (\underline{aa_\bullet^2})^{-1} (\underline{aa_\bullet})^{-1}. \end{aligned}$$

Therefore the reduced characteristic sequence of $ABACBC$ is

$$1, a_\bullet, -aa_\bullet, -1, a, aa_\bullet, -a^2 a_\bullet, aa_\bullet, a^2 a_\bullet^2, -aa_\bullet^2, -aa_\bullet.$$

It differs from the one of $ACAC$ already in the first term. Therefore $ABACBC$ is not homotopic to $ACAC$. As a check of our computations, note that the sum of all terms in the characteristic sequences of these two nanowords is the same, as it should be because these nanowords are indistinguishable by the invariant λ .

REFERENCES

- [FR] R. Fenn, C. Rourke, *Racks and links in codimension two*, J. Knot Theory Ramifications 1 (1992), no. 4, 343–406.
- [Ka] S. Kamada, *Knot invariants derived from quandles and racks*. Invariants of knots and 3-manifolds (Kyoto, 2001), 103–117 (electronic), Geom. Topol. Monogr., 4, Geom. Topol. Publ., Coventry, 2002.

- [Ro] P. Rosenstiehl, *Solution algébrique du problème de Gauss sur la permutation des points d'intersection d'une ou plusieurs courbes fermées du plan*, C. R. Acad. Sci. Paris Sér. A-B 283 (1976), no. 8, A551–A553.
- [SW] D. Silver, S. Williams, *An invariant for open virtual strings*, math.GT/0409185.
- [Tu1] V. Turaev, *Virtual strings*, math.GT/0311185, to appear in Ann. Inst. Fourier.

IRMA, UNIVERSITÉ LOUIS PASTEUR - C.N.R.S.,
 7 RUE RENÉ DESCARTES
 F-67084 STRASBOURG
 FRANCE
 E-MAIL: TURAEV@MATH.U-STRASBG.FR